

東邦大学学術リポジトリ

Toho University Academic Repository

| | |
|-----------|---|
| タイトル | Retrospective study of colposcopy for precancerous cervical lesions and correlation between diagnosis and number of biopsy specimens |
| 別タイトル | 子宮頸部前癌病変に対するコルポスコピーと、診断と生検個数に関する後方視的検討 |
| 作成者（著者） | 高橋, 怜奈 |
| 公開者 | 東邦大学 |
| 発行日 | 2020.03.15 |
| 掲載情報 | 東邦大学大学院医学研究科 博士論文. |
| 資料種別 | 学位論文 |
| 内容記述 | 主査：澁谷和俊 / タイトル：Retrospective study of colposcopy for precancerous cervical lesions and correlation between diagnosis and number of biopsy specimens / 著者：Rena Takahashi, Kaneyuki Kubushiro, Isao Murakami, Tsuyoki Kugimiya, Kyoko Tanaka, Mineto Morita / 掲載誌：Toho Journal of Medicine / 巻号・発行年等：6(3):99-103, 2020 / 本文ファイル: 出版社版 |
| 著者版フラグ | ETD |
| 報告番号 | 32661甲第954号 |
| 学位記番号 | 甲第659号 |
| 学位授与年月日 | 2020.03.15 |
| 学位授与機関 | 東邦大学 |
| DOI | info:10.14994/tohojmed.2019_025 |
| その他資源識別子 | info:10.14994/tohojmed.2019_025 |
| メタデータのURL | https://mylibrary.toho-u.ac.jp/webopac/TD11304270 |

Retrospective Study of Colposcopy for Precancerous Cervical Lesions and Correlation between Diagnosis and Number of Biopsy Specimens

Rena Takahashi^{1,2)} Kaneyuki Kubushiro³⁾ Isao Murakami¹⁾
Tsuyoki Kugimiya¹⁾ Kyoko Tanaka¹⁾ and Mineto Morita^{4)*}

¹⁾Department of Gynecology, Toho University Ohashi Medical Center, Tokyo, Japan

²⁾Department of Obstetrics and Gynecology, Toho University Graduate School of Medicine, Tokyo, Japan

³⁾Tokyo Health Service Association, Tokyo, Japan

⁴⁾Department of Obstetrics and Gynecology, Toho University Omori Medical Center, Tokyo, Japan

ABSTRACT

Introduction: Histological diagnosis is performed in patients with abnormal findings on cervical cytology and macroscopic abnormalities of the cervix. In general, targeted histological diagnosis is performed to extract the specimens from lesions with abnormal findings under a colposcope. In this study, we aim to retrospectively verify the number of specimens from different cervical lesions that are most effective for diagnosis.

Methods: We included women with abnormal findings on cervical cytology who underwent targeted biopsy under a colposcope in our hospital from August 2013 to March 2016. The number of specimens extracted and the number of biopsies required to achieve diagnosis were examined in patients with cervical histopathology showing cervical intraepithelial neoplasia (CIN)1, CIN2, and CIN3. Furthermore, when cervical histology showed indications for conization and total hysterectomy, these procedures were performed to confirm the ultimate histopathological diagnosis.

Results: The definite diagnosis of CIN was made from one, one and two, and one, two, and three biopsies in 81.6%, 96.1%, and 100% patients, respectively.

Conclusions: All patients with CIN1 to CIN3 were diagnosed with three biopsies. In particular, higher number of biopsies was associated with higher rates of definite diagnosis. However, in no cases, definite CIN diagnosis was made at the four biopsies, suggesting that four or more cervical histological biopsies are generally unnecessary. Furthermore, given that 100% patients were diagnosed by their three biopsies, and at most, only three biopsies should be performed.

Toho J Med 6 (3): 99–103, 2020

KEYWORDS: colposcopy, cervical intraepithelial neoplasia, biopsy, cervical cancer

*Corresponding Author: Mineto Morita, 6-11-1 Omorinishi, Ota, Tokyo 143-8541, Japan, tel: +81 3 3762 4151
e-mail: mmorita@med.toho-u.ac.jp
DOI: 10.14994/tohojmed.2019-025

Received Oct. 31, 2019; Accepted Dec. 19, 2019
Toho Journal of Medicine 6 (3), Sept. 1, 2020.
ISSN 2189-1990, CODEN: TJMOA2

Introduction

Cervical cancer is the fourth most common cancer in women with an estimated 570,000 new cases in 2018 representing 7.5% of all female cancer death.¹⁾ Cervical cancer is the easiest gynecologic cancer to prevent with regular screening tests and follow-up; it is highly curable when found and treated early.²⁾

The detection of cytological abnormalities through screening has resulted in a fall in the incidence of cervical cancer because of the treatment of these pre-invasive lesions.³⁾

Colposcopic examination of the cervix allows assessment of the abnormality before it is treated, either by excision or ablation.⁴⁻⁶⁾ Colposcopy is a subjective test; its purpose is to identify disease, obtain representative specimens for histology, and direct patient management.^{7,8)}

In 2013, the American Society for Colposcopy and Cervical Pathology updated its guidelines to suggest that a histological diagnosis should be done for patients with abnormalities on cervical cytology or those with macroscopic abnormalities of the cervix.⁹⁾ The punch biopsy also plays a role in the management of women undergoing ablative treatment for CIN because pretreatment biopsies are required to exclude invasive disease.^{10,11)} Generally, targeted histological biopsy is performed to directly extract specimens from the lesions with abnormal findings under a colposcope. However, colposcopy is a subjective test, and its accuracy has been recently questioned.^{12,13)} According to Nakamura et al., CIN2 and more severe lesions are diagnosed by one-lesion biopsy in 78.1% patients and by two-lesion biopsies in $\geq 90\%$, regardless of the tester.¹⁴⁾ Stoler MH et al. reported that one, two, and three or more biopsies led to 47%, 65%, and 77% of definite diagnoses, respectively. Collecting specimens from more lesions is historically recommended to improve the accuracy of cervical tissue biopsy.¹⁵⁾ However, cervical histological diagnosis, an invasive test, causes bleeding, and it is ideal to establish a certain diagnosis with a less-invasive one.

Therefore, we investigated the number of biopsies required for accurate histological diagnosis in 350 patients who underwent colposcopy and histological biopsy. In those requiring conization, histological diagnoses before and after conization were also compared.

Methods

We included women with abnormal findings on cervical

cytology who underwent colposcopy and histological biopsy from August 2013 to March 2016 at our hospital. Nine testers were involved. We investigated the number of biopsies and their lesions required for the cervical histopathological diagnoses of CIN1, CIN2, and CIN3. Furthermore, the preoperative histological and histopathological diagnoses of the conization specimens in patients who underwent conization or total hysterectomy were analyzed. The order of biopsy was decided as per the tester's discretion. Hematoxylin and eosin staining and the World Health Organization classifications were used for histopathological diagnosis. Moreover, when cervical histology showed indications for conization and total hysterectomy, these procedures were performed to confirm the ultimate histopathological diagnosis.

This study was approved by the relevant ethics committee of Toho University Ohashi Medical Center (No. H 17018).

Results

A total of 350 female patients were histologically diagnosed with chronic cervicitis, CIN, and adenocarcinoma in situ (AIS). Their characteristics are shown in Table 1. In terms of age, 71 (20.3%), 137(39.1%), 86 (24.6%), and 56 (16.0%) were in their 20s, 30s, 40s, and 50s, respectively. The median age was 36 (20-79) years. In addition, the results of cervical cytology indicated ASC-US, LSIL, HSIL, AGC, ASC-H, CIS, and SCC in 24 (6.9%), 133 (46.6%), 177 (50.6%), 5 (1.4%), 3 (0.86%), 3 (0.86%), and 5 (1.4%) patients, respectively.

Targeted histological diagnosis under a colposcope revealed chronic cervicitis, CIN1, CIN2, CIN3, and AIS in 65 (18.6%), 107 (30.6%), 89 (25.4%), 87 (24.9%), and 2 (0.57%) patients, respectively. Furthermore, 43 (12.3%), 153 (43.7%), 132 (37.7%), and 22 (6.3%) patients underwent one, two, three, and four biopsies, respectively. As shown in Table 2, 231 (81.6%), 272 (96.1%), and 283 (100%) of them were diagnosed at their one, one and two, and one, two, and three biopsies, respectively.

Furthermore, we investigated the number of biopsies by each CIN grade and the number of biopsies required for diagnosis. In CIN1, one, two, three, and four biopsies were taken in 14 (13.1%), 58 (54.2%), 32 (29.9%), and 3 (2.8%) patients, respectively. Furthermore, 89 (83.2%), 105 (98.1%), and 107 (100%) patients were diagnosed with CIN1 at their one, one and two, and one, two, and three biopsies, respectively (Fig. 1). In CIN2, one, two, three, and four biopsies

Table 1 Characteristics of the study participants

| Characteristics | N | % |
|--|-----|------|
| Age (years) | | |
| 20-29 | 71 | 20.3 |
| 30-39 | 137 | 39.1 |
| 40-49 | 86 | 24.6 |
| ≥ 50 | 56 | 16.0 |
| Cytology | | |
| ASC-US | 24 | 6.9 |
| LSIL | 133 | 46.6 |
| ASC-H | 3 | 0.86 |
| HSIL | 177 | 50.6 |
| AGC | 5 | 1.4 |
| CIS | 3 | 0.86 |
| SCC | 5 | 1.4 |
| Histology | | |
| Total | 350 | 100 |
| Chronic cervicitis | 65 | 18.6 |
| CIN1 | 107 | 30.6 |
| CIN2 | 89 | 25.4 |
| CIN3 | 87 | 24.9 |
| AIS | 2 | 0.57 |
| Number of cervical specimens (CIN1, CIN2, CIN3) | | |
| Total | 283 | 100 |
| One | 29 | 10.2 |
| Two | 117 | 41.3 |
| Three | 120 | 42.4 |
| Four | 17 | 6.0 |
| Number of biopsies to diagnosis (CIN1, CIN2, CIN3) | | |
| Total | 283 | 100 |
| One | 231 | 81.6 |
| One and two | 272 | 96.1 |
| One, two, and three | 283 | 100 |

were taken in 9 (10.1%), 36 (40.4%), 41 (46.1%), and 3 (3.4%) patients, respectively. Furthermore, CIN2 diagnoses were reached at the one, one and two, and one, two, and three biopsies in 75 (84.3%), 87 (97.8%), 89 (100%) patients, respectively (Fig. 2). In CIN3, one, two, three, and four biopsies were performed in 6 (6.9%), 23 (26.4%), 47 (54.0%), and 11 (12.6%) patients, respectively. Furthermore, 67 (77.0%), 80 (83.8%), and 87 (100%) patients were diagnosed with CIN3 at their one, one and two, and one, two, and three biopsies, respectively (Fig. 3). Thus, CIN diagnosis was established at the one, one and two, and one, two, and three biopsies in 81.6%, 96.1%, and 100% patients, respectively.

A total of 81 CIN3 patients underwent cervical conization or total hysterectomy, of which 59, 8, and 14 patients were diagnosed with CIN3, CIN2, and chronic cervicitis by histopathology, respectively. Furthermore, of the five CIN

Table 2 Number of cervical specimens and number of biopsies to diagnose (CIN)

| Number of cervical specimens and number of biopsies to diagnose (CIN1, CIN2, and CIN3) | N | % |
|--|-----|------|
| Number of cervical specimens (CIN1) | | |
| Total | 107 | 100 |
| One | 14 | 13.1 |
| Two | 58 | 54.2 |
| Three | 32 | 29.9 |
| Four | 3 | 2.8 |
| Number of biopsies to diagnose (CIN1) | | |
| Total | 107 | 100 |
| One | 89 | 83.2 |
| One and two | 105 | 98.1 |
| One, two, and three | 107 | 100 |
| Number of cervical specimens (CIN2) | | |
| Total | 89 | 100 |
| One | 9 | 10.1 |
| Two | 36 | 40.4 |
| Three | 41 | 46.1 |
| Four | 3 | 3.4 |
| Number of biopsies to diagnose (CIN2) | | |
| Total | 89 | 100 |
| One | 75 | 84.3 |
| One and two | 87 | 97.8 |
| One, two, and three | 89 | 100 |
| Number of cervical specimens (CIN3) | | |
| Total | 87 | 100 |
| One | 6 | 6.9 |
| Two | 23 | 26.4 |
| Three | 47 | 54.0 |
| Four | 11 | 12.6 |
| Number of biopsies to diagnose (CIN3) | | |
| Total | 87 | 100 |
| One | 67 | 77.0 |
| One and two | 80 | 92.0 |
| One, two, and three | 87 | 100 |

2 patients who underwent this procedure, four patients were diagnosed with CIN3, and one patient was diagnosed with CIN2, histopathologically.

Discussion

We studied 350 patients with abnormal findings in cervical cytology. All were diagnosed with CIN1-CIN3 by the three biopsies. Furthermore, with more number of biopsies, definite diagnoses were made at a higher probability. However, no patient was definitely diagnosed at the four biopsies. Therefore, four or more biopsies are generally considered unnecessary for cervical histology. In addition, given that 100% patients were diagnosed by their three bi-




Fig. 1

Number of biopsies and detection rates of CIN1 diagnosis
Overall 83.2% patients were diagnosed at one biopsy, 98.1% were diagnosed at one and two biopsies, and 100% were diagnosed at one, two, and three biopsies.




Fig. 3

Number of biopsies and detection rates of CIN3 diagnosis
Overall 77.0% patients were diagnosed at one biopsy, 92.0% were diagnosed at one and two biopsies, and 100% were diagnosed at one, two, and three biopsies.




Fig. 2

Number of biopsies and detection rates of CIN2 diagnosis
Overall 84.3% patients were diagnosed at one biopsy, 97.8% were diagnosed at one and two biopsies, and 100% were diagnosed at one, two, and three biopsies.

Table 3 Association between CIN and the sensitivity of colposcopic procedure, Chi-square test

| | one | two | three | |
|------|-----|-----|-------|-----|
| CIN1 | 89 | 16 | 2 | 107 |
| CIN2 | 75 | 12 | 2 | 89 |
| CIN3 | 67 | 13 | 7 | 87 |
| | 231 | 41 | 11 | 283 |

P-value = 0.1967

opies, and at most, only three biopsies should be performed.

Furthermore, the results of cervical histology after conization or total hysterectomy were investigated. The most severe abnormality found in the present study was CIN3.

In addition to the detection rate of CIN3 is seem lower than CIN1 and CIN2, there is no significant difference (Table 3).

However, future studies should investigate more number of cases to cognize the reason.

There are inconsistent results regarding the detection rates of colposcopic diagnosis.

Nicolas et al. reported that HSIL was detected in biopsies of the specimens extracted from one, two, and three

lesions in 60.6%, 85.6%, and 95.6% patients, respectively.¹⁶⁾

According to Nakamura et al., 78.1% CIN2 or severe lesions were detected at the first biopsy, but at least two biopsies should be taken.¹⁴⁾ In contrast, Robert et al. showed that CIN2 and severe lesions were colposcopically diagnosed in only 57.1% patients.¹⁷⁾ Because the detection rates differ between these reports, there is no consensus on the number of biopsies required for cervical histological diagnosis.

We considered that the difference of detection rates occurs from difference from examiners of each hospital. In this study, nine examiners were involved, but some of them were enrolled in a short period of time and nine examiners aren't corresponded to six authors listed. The order of biopsy is different depending on the examiners; therefore, to obtain more accurate data, a number of examiners should be limited or should unify the order of biopsy when there are more than one examiner.

There are several limitations of the study. First, given that the order of specimen extractions varied as per the

tester who performed the cervical histology, the accurate number of biopsies required for definite diagnosis has not been necessarily clarified. For example, in performing a biopsy of the upper and lower lesions of the cervix, in some cases, even if the upper lesion was stronger, the specimen is extracted first from the lower lesion to prevent the invasion of blood flowing from the biopsy site of the upper lesion. However, biopsies from four sites are considered unnecessary.

To obtain more accurate data, future studies should standardize the order of cervical biopsies between testers from the most severe lesions to the less severe ones.

Conflicts of interest: None declared.

References

- 1) Human papillomavirus (HPV) and cervical cancer (2019). [Accessed 3 March 2019] [https://www.who.int/en/news-room/fact-sheets/detail/human-papillomavirus-\(hpv\)-and-cervical-cancer](https://www.who.int/en/news-room/fact-sheets/detail/human-papillomavirus-(hpv)-and-cervical-cancer)
- 2) Inside knowledge: get the facts about gynecologic cancer (2014). https://www.cdc.gov/cancer/knowledge/publications/fact_sheets.htm
- 3) Sasieni P, Adams J. Effect of screening on cervical cancer mortality in England and Wales: analysis of trends with an age period cohort model. *BMJ*. 1999; 318: 1244-5.
- 4) NHSCSP. Colposcopy and Programme Management: Guidelines for the NHS Cervical Screening Programme. 2nd ed. London: NHSCSP; 2010.
- 5) Jordan J, Arbyn M, Martin-Hirsch P, Schenck U, Baldauf J-J, Da Silva D, et al. European guidelines for quality assurance in cervical cancer screening: recommendations for clinical management of abnormal cervical cytology, part 1. *Cytopathology*. 2008; 19: 342-54.
- 6) Arbyn M, Anttila A, Jordan J, Ronco G, Schenck U, Segnan N, et al. European guidelines for quality assurance in cervical cancer screening. Second edition - summary document. *Ann Oncol*. 2010; 21: 448-58.
- 7) Jeronimo J, Massad LS, Castle PE, Wacholder S, Schiffman M. Interobserver agreement in the evaluation of digitized cervical images. *Obstet Gynecol*. 2007; 110: 833-40.
- 8) Apgar B, Brotzman G, Spitzer M, Ignatavicius DD. In: Apgar B, Brotzman G, Spitzer M, Ignatavicius DD, editors. *Colposcopy, principles and practice: an integrated textbook and atlas*. 1st ed. Philadelphia, PA: WB Saunders; 2002. p. 463-71.
- 9) Massad L, Einstein MH, Huh WK, Katki HA, Kinney WK, Schiffman MS. 2012 ASCCP Consensus Guidelines Conference. Updated consensus guidelines for the management of abnormal cervical cancer screening tests and cancer precursors. *J Obstet Gynaecol Can*. 2012; 34: 1188.
- 10) Anderson MC. Invasive carcinoma of the cervix following local destructive treatment for cervical intraepithelial neoplasia. *Br J Obstet Gynaecol*. 1993; 100: 657-63.
- 11) Shumsky AG, Stuart GC, Nation J. Carcinoma of the cervix following conservative management of cervical intraepithelial neoplasia. *Gynecol Oncol*. 1994; 53: 50-4.
- 12) Jeronimo J, Schiffman M. Colposcopy at a crossroads. *Am J Obstet Gynecol*. 2006; 195: 349-53.
- 13) Chase DM, Kalouyan M, DiSaia PJ. Colposcopy to evaluate abnormal cervical cytology in 2008. *Am J Obstet Gynecol*. 2009; 200: 472-80.
- 14) Nakamura Y. Optimizing biopsy procedures during colposcopy for women with abnormal cervical cancer screening results: a multicenter prospective study. *Int J Clin Oncol*. 2015; 20: 579-85.
- 15) Stoler MH. The accuracy of colposcopic biopsy: analyzes from the placebo arm of the Gardasil clinical trials. *Int J Cancer*. 2011; 128: 1354-62.
- 16) Wentzensen N, Walker JL, Gold MA, Smith KM, Zuna RE, Mathews C, et al. Multiple biopsies and detection of cervical cancer precursors at colposcopy. *J Clin Oncol*. 2015; 33: 83.
- 17) Pretorius RG, Zhang WH, Belinson JL, Huang MN, Wu LY, Zhang X, et al. Colposcopically directed biopsy, random cervical biopsy, and endocervical curettage in the diagnosis of cervical intraepithelial neoplasia II or worse. *Am J Obstet Gynecol*. 2004; 191: 430-4.

©Medical Society of Toho University. Toho Journal of Medicine is an Open Access journal distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view the details of this license, please visit (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).