

東邦大学学術リポジトリ



OPAC

東邦大学メディアセンター

タイトル	Asian dust and pediatric emergency department visits due to bronchial asthma and respiratory diseases in Nagasaki, Japan
別タイトル	長崎市における黄砂と気管支喘息や呼吸器疾患による小児救急受診との関連
作成者(著者)	中村, 孝裕
公開者	東邦大学
発行日	2016.03
掲載情報	東邦大学大学院医学研究科 博士論文. 11.
資料種別	学位論文
内容記述	主査: 村上義孝 / タイトル: Asian dust and pediatric emergency department visits due to bronchial asthma and respiratory diseases in Nagasaki, Japan / 著者: Takahiro Nakamura, Masahiro Hashizume, Kayo Ueda, Atsushi Shimizu, Ayano Takeuchi, Tatsuhiko Kubo, Kunio Hashimoto, Hiroyuki Moriuchi, Hiroshi Odajima, Tasuku Kitajima, Kasumi Tashiro, Kunio Tomimasu, Yuji Nishiwaki / 掲載誌: Journal of Epidemiology / 巻号・発行年等: 26(11):593-601, 2016 / 本文ファイル: 出版者版
著者版フラグ	ETD
報告番号	32661甲第821号
学位記番号	甲第552号
学位授与年月日	2017.03.28
学位授与機関	東邦大学
DOI	info:doi/10.2188/jea.JE20150309
その他資源識別子	https://www.jstage.jst.go.jp/article/jea/26/11/26_JE20150309/article/char/ja/
メタデータのURL	https://mylibrary.toho-u.ac.jp/webopac/TD20910995



Asian Dust and Pediatric Emergency Department Visits Due to Bronchial Asthma and Respiratory Diseases in Nagasaki, Japan

Takahiro Nakamura¹, Masahiro Hashizume², Kayo Ueda³, Atsushi Shimizu⁴, Ayano Takeuchi⁵, Tatsuhiko Kubo⁶, Kunio Hashimoto⁷, Hiroyuki Moriuchi⁷, Hiroshi Odajima⁸, Tasuku Kitajima⁹, Kasumi Tashiro¹⁰, Kunio Tomimasu¹¹, and Yuji Nishiwaki¹

¹Department of Environmental and Occupational Health, School of Medicine, Toho University, Tokyo, Japan

²Department of Paediatric Infectious Diseases, Institute of Tropical Medicine, Nagasaki University, Nagasaki, Japan

³Department of Environmental Engineering, Kyoto University Graduate School of Engineering, Kyoto, Japan

⁴Regional Atmospheric Environment Section Center, National Institute for Environmental Studies, Tsukuba, Japan

⁵Department of Preventive Medicine and Public Health, School of Medicine, Keio University, Tokyo, Japan

⁶Department of Public Health, School of Medicine, University of Occupational and Environmental Health, Kitakyushu, Fukuoka, Japan

⁷Department of Paediatrics, Graduate School of Biomedical Sciences, Nagasaki University, Nagasaki, Japan

⁸Department of Paediatrics, National Hospital Organization Fukuoka Hospital, Fukuoka, Japan

⁹Department of Paediatrics, Kamigoto Hospital, Nagasaki, Japan

¹⁰Department of Paediatrics, Isahaya General Hospital, Nagasaki, Japan

¹¹Nagasaki Municipal Primary Emergency Medical Center, Nagasaki, Japan

Received November 11, 2015; accepted December 29, 2015; released online May 14, 2016

Copyright © 2016 Takahiro Nakamura et al. This is an open access article distributed under the terms of Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

ABSTRACT

Background: The adverse health effects of Asian dust (AD) on the respiratory system of children are unclear. We hypothesized that AD events may lead to increased visits by children to emergency medical centers due to bronchial asthma and respiratory diseases, including bronchial asthma.

Methods: We used anonymized data on children receiving primary emergency treatment at Nagasaki Municipal Primary Emergency Medical Center, Japan between March 2010 and September 2013. We used Light Detection and Ranging (LIDAR) data to assess AD exposure and performed time-stratified case-crossover analyses to examine the association between AD exposure and emergency department visits. The main analysis was done with data collected from March through May each year.

Results: The total number of emergency department visits during the study period was 756 for bronchial asthma and 5421 for respiratory diseases, and the number of “AD days” was 47. In school children, AD events at lag day 3 and lag day 4 were associated with increased emergency department visits due to bronchial asthma, with odds ratios of 1.837 (95% confidence interval [CI], 1.212–2.786) and 1.829 (95% CI, 1.179–2.806), respectively. AD events were significantly associated with respiratory diseases among preschool children at lag day 0, lag day 1, and lag day 2, with odds ratios of 1.244 (95% CI, 1.128–1.373), 1.314 (95% CI, 1.189–1.452), and 1.273 (95% CI, 1.152–1.408), respectively. These associations were also significant when the results were adjusted for meteorological variables and other air pollutants.

Conclusions: The study findings suggested that AD exposure increases emergency department visits by children.

Key words: Asian dust; emergency department visits; bronchial asthma; respiratory diseases; children

INTRODUCTION

Asian dust (AD) is a natural phenomenon in which wind carries dust over large distances from the Yellow River basin and deserts in northern China and Mongolia. It is now regarded

as a worldwide environmental problem caused by manmade deforestation, soil degradation, and desertification.^{1–3}

AD reaches Japan throughout the year, but the amount increases from February and peaks in April.⁴ AD is a common problem throughout Northeast Asia, but the type and degree of

Address for correspondence: Yuji Nishiwaki, MD, MSc, DMSc, Department of Environmental and Occupational Health, School of Medicine, Toho University, 5-21-16 Omori-Nishi, Ota-ku, Tokyo 143-8540, Japan (e-mail: yuji.nishiwaki@med.toho-u.ac.jp).




Figure 1. Map of Japan (left) and Nagasaki City (right)
 ○ shows Nagasaki Prefecture (left); ◇ shows Nagasaki Municipal Primary Emergency Medical Center (right);
 ◆ shows LIDAR in Omura City (right); ● shows four air pollution monitoring stations in Nagasaki city (right);
 ▲ shows Nagasaki Marine Observatory.
 LIDAR, light detection and ranging.

damage it causes may vary, depending on the distance from the source of the dust.¹ In Japan, AD most often affects western Japan. Nagasaki Municipal Primary Emergency Medical (NMPEM) Center is located in the center of the city and is the primary night-time and holiday emergency hospital for children living in the Nagasaki area. Children appear to be more vulnerable than adults, in terms of both susceptibility and response to environmental changes, including air pollution.⁵⁻⁷

Although studies on the effects of AD on the respiratory system in children have been carried out in Taiwan, China, and Korea,⁸⁻¹³ the results have been contradictory. Some studies reported an increased risk of clinic visits⁸ and hospitalization for asthma,¹⁴ whereas another study reported a negative association with emergency asthma hospitalization.¹⁵

We speculated that the effects of AD may vary between preschool and school children because of differences in their daily activities. However, there is very little evidence to support this. In addition, if AD does indeed affect the respiratory system of children, the dose-response relationship needs to be investigated. Recent epidemiological studies from Korea and Japan have measured levels of AD using Light Detection and Ranging (LIDAR).^{14,16,17} LIDAR is a type of radar that uses laser beams instead of radio waves. The laser beams emitted from the ground are scattered by fine particles in the air, and, by measuring the scattered laser light, operators can determine the vertical distribution of dust particle concentration and changes over time.¹⁸ Such measurements provide reliable quantitative data on AD levels in the atmosphere.

The objective of this study was to examine data from the NMPEM Center to identify any relationship between AD and emergency department visits by children.

METHODS

Study area

Nagasaki Prefecture borders Saga Prefecture to the east, but it is otherwise surrounded by the East China Sea (Figure 1). The prefecture's capital is Nagasaki City. The NMPEM Center serves the whole of Nagasaki City and two neighboring towns (Togitsu and Nagayo in Nishisonogi District). In 2013, the total population of this area was approximately 500 000, with about 25 000 children aged 0-5 years and about 46 000 children aged 6-15 years.¹⁹ Approximately 90% of the patients who access the medical center live in the above-mentioned area, according to unpublished information from the center. The center offers emergency services for children 7 days per week from 20:00 to 06:00. Several pediatricians work in shifts to provide these services.

Collection of data on emergency department visits

We obtained unlinkable anonymized data (research ID number date of visits, date of birth, diagnosis, and treatment) on all children aged 0-15 years who received primary emergency treatment at the Center between March 2010 and September 2013. We created two diagnostic categories: 1) bronchial asthma and 2) respiratory diseases, including bronchial asthma, bronchitis, pharyngitis, and common cold.

In order to focus on the effects of AD, we excluded patients diagnosed with specific respiratory pathogens, such as influenza viruses, respiratory syncytial virus, or *mycoplasma* infections.

Measurement of AD

To assess exposure to AD during the study period, we used LIDAR data obtained from an installation in Omura City (Figure 1), which is approximately 30 km from Nagasaki City. LIDAR utilizes polarized laser light to recognize shape differences and can distinguish AD particles from other air pollutants, which are generally spherical. If the lower atmosphere is well mixed, the concentration of AD on the ground is similar to that between 120 m and 270 m above ground, the lowest level for which LIDAR data are available.¹⁸ The National Institute for Environmental Studies LIDAR network can estimate the dust extinction coefficients of non-spherical and spherical components.²⁰ LIDAR measurements were compiled as total 24-hour data (midnight to midnight). AD extinction coefficients larger than 1/km and spherical extinction coefficients larger than 2/km were excluded, because these data may be influenced by meteorological factors, such as cloud, fog, rain, and snow.²⁰

In our study, we defined “AD days” as when all of the following criteria were met²¹:

1. The daily maximum suspended particulate matter (SPM) concentration was over 50 $\mu\text{g}/\text{m}^3$.
2. The daily maximum AD extinction coefficient as measured by LIDAR was more than 0.05/km.
3. The correlation coefficient between the hourly AD extinction coefficient and hourly SPM concentration was a fixed value.

For SPM concentrations, we used the daily average of hourly measurements made by four air pollution monitoring stations. SPM is defined under the Japanese Air Quality Standard as any particle with an upper 100% cut-off point of 10 μm in aerodynamic diameter.²²

Other air pollutants and meteorological data

Data on other air pollutants (sulfur dioxide [SO_2], nitrogen dioxide [NO_2], and photochemical oxidants [Ox]) were collected from four air pollution monitoring stations in Nagasaki City. Daily average concentrations of SO_2 and NO_2 were calculated from hourly concentrations measured at each station. Ox is defined as mixtures of ozone and other secondary oxidants generated by photochemical reactions and is considered to be a proxy for ozone. Maximum hourly concentrations of Ox measured at each station were used. Data based on fewer than 20 hourly measurements on any one day were excluded.^{15,20}

Daily temperature and relative humidity data in Nagasaki City were obtained from the Nagasaki Marine Observatory of the Japanese Meteorological Agency. We calculated the daily average temperature and relative humidity from data collected

hourly at Nagasaki Marine Observatory (between 00:00 and 23:00), and recorded the maximum and minimum temperatures.

Statistical analysis

We used the Mann-Whitney U test or Student's *t*-tests to compare AD extinction coefficients, spherical extinction coefficients, SPM, other air pollutants and meteorological data on AD days and non-AD days.

A time-stratified case-crossover analysis was used to examine the relationship between AD and emergency department visits due to bronchial asthma and respiratory diseases. In the same way as in a matched case-control analysis, this analysis assigned the day on which a patient visited the clinic with bronchial asthma or respiratory disease as the case day, and comparisons were made with control days chosen on the same day of the week earlier or later in the same month of the same year.^{23,24} The advantage of this design is that it controls for time-invariant personal factors, the effects of long-term trends, seasonality, and the day of the week. The strength of the association between AD days and emergency clinic visits was shown by the odds ratios (ORs) and 95% confidence intervals (CIs) of the conditional logistic models. *P* values of less than 0.05 were considered to indicate statistical significance.

Following the Japanese Pediatric Guideline for the Treatment and Management of Asthma 2012, we classified three age categories: under 2 years old, 2 through 5 years old, and 6 through 15 years old²⁵; however, because we found that the results for the first two groups were similar, we combined them in the subsequent analysis. Thus, the preschool children group consisted of patients aged 0 through 5 years, and the school children group consisted of those aged 6 through 15 years. We analyzed combined data for boys and girls, because the associations between AD and emergency department visits did not differ by sex.

In addition to AD days, we added temperature and relative humidity as co-variables,²⁰ along with SO_2 , NO_2 , and Ox concentrations, and the spherical extinction coefficients (these variables were average on the case day, lag day 1, and lag day 2). We constructed three adjusted models. In the basic model, we adjusted for temperature and relative humidity. In the single-pollutant model, we adjusted for one of the air pollutants (SO_2 , NO_2 , or Ox), in addition to the variables in the basic model. In the multi-pollutant model, we adjusted for two of the air pollutants, in addition to the variables in the basic model.

Because the effects of AD can persist over several days, we examined the effects with several lag times; the case day (lag day 0), lag day 1, lag day 2, lag day 3, lag day 4, and lag day 5 as a single lag model (the model included only one exposure variable). We also examined the lagged cumulative effects from lag day 0 to day 1 (designated lag 01), lag day 0 to day 2 (lag 02), lag day 0 to day 3 (lag 03), lag day 0 to day 4 (lag 04), and lag day 0 to day 5 (lag 05).




Figure 2. Number of Asian dust days from March 2010 through September 2013.

In the basic model, we used not only the binomial index of AD but also the AD extinction coefficients, with reference to the methods of Ueda et al.²⁶ We classified results into three categories: Non-AD days (extinction coefficient $<0.047/\text{km}$), moderate AD days (extinction coefficient $0.047\text{--}0.065/\text{km}$), and heavy AD days (extinction coefficient $\geq 0.066/\text{km}$), according to the median ($0.047/\text{km}$) and 75th percentile ($0.066/\text{km}$) value of the AD extinction coefficients.

In addition, several sensitivity analyses were performed. First, because airway inflammation and sensitivity to environmental stimuli may persist after an initial attack, we carried out further analyses after excluding visits within 4 weeks of the first visit. Second, we conducted analyses using data for the whole year rather than just those for March through May.

All analyses were performed with STATA ver. 12.1 (StataCorp, College Station, TX, USA).

Ethical considerations

This study was a retrospective observational study using unlinkable anonymized data. It was approved by the Ethics Committee of Nagasaki University Hospital (Approval No. 13082663) and Toho University School of Medicine (Approval No. 25100).

RESULTS

AD days

LIDAR measurements showed that the total number of AD days during the study period (March 2010 through September 2013) was 63 (Figure 2). The largest number of AD days was observed in 2010, and most (47/63) occurred in spring (March to May) each year, although they were also occasionally observed in November and December. Therefore, most of our analyses were based on data collected in March through May.

Primary emergency department visits to Nagasaki Municipal Primary Emergency Medical Center

During the 3.5-year study period (March 2010 through September 2013) there were 43 892 emergency department visits, of which 34 170 were made by patients aged 15 years or below (18 805 boys and 15 365 girls). Of these, 9937 were made between March and May; 756 involved bronchial asthma (mean age of the patients, 3.78 years) and 5421 were due to respiratory diseases (mean age, 3.45 years) (Table 1). Boys and preschool children made up the majority in both diagnostic categories. The largest number of visits due to bronchial asthma occurred in May 2010, and the smallest in March 2012; May 2011 saw the largest number of visits due to respiratory diseases, and March 2010 saw the smallest.

AD, other air pollutants, and meteorological data

We compared the levels of AD, other air pollutants, and meteorological data on AD days and non-AD days from March through May, 2010–2013 (Table 2). The mean AD coefficient on AD days ($0.07/\text{km}$) was significantly higher than that on non-AD days ($0.03/\text{km}$). There were differences in the concentrations of other air pollutants and meteorological variables (except NO_2 and relative humidity) on AD days and non-AD days.

Association between AD and emergency department visits

The crude ORs by age group for emergency department visits on AD days due to bronchial asthma and respiratory diseases are shown in Figure 3.

In all children, AD events at lag day 2, lag day 3, and lag day 4 were associated with increased emergency department visits due to bronchial asthma, with ORs of 1.343 (95% CI,

Table 1. Number of emergency department visits (March to May)

		Bronchial asthma			Respiratory diseases		
		All Children 0–15 years old (n = 756)	Preschool Children 0–5 years old (n = 551)	School Children 6–15 years old (n = 205)	All Children 0–15 years old (n = 5421)	Preschool Children 0–5 years old (n = 4070)	School Children 6–15 years old (n = 1351)
Age, years	Mean	3.78	1.91	8.82	3.45	1.5	9.35
	Median	3	2	9	2	1	9
Sex	Male	474	337	137	2942	2198	744
	Female	282	214	68	2479	1872	607
Year	2010	214	159	55	1205	975	230
	2011	197	138	59	1485	1111	374
	2012	156	112	44	1343	956	387
	2013	189	142	47	1388	1028	360
Month	March	212	154	58	1626	1150	476
	April	250	182	68	1749	1340	409
	May	294	215	79	2046	1580	466

Table 2. Summary of data on Asian dust, other air pollutants, and meteorological factors (March 2010–May 2013)

	Non-Asian-dust days						Asian-dust days						P value ^{a,b}
	Number of days	Mean	SD	25th percentile	Median	75th percentile	Number of days	Mean	SD	25th percentile	Median	75th percentile	
Asian dust extinction coefficients, /km	320	0.03	0.03	0.01	0.02	0.03	47	0.07	0.08	0.03	0.05	0.07	<0.01 ^a
Spherical extinction coefficients, /km	320	0.13	0.11	0.05	0.10	0.16	47	0.19	0.13	0.09	0.16	0.24	<0.01 ^a
SO ₂ , ppb	321	1.52	0.77	1.00	1.30	1.80	47	1.98	0.97	1.20	1.60	2.50	<0.01 ^a
NO ₂ , ppb	321	7.58	2.28	6.00	7.40	9.00	47	8.11	2.60	6.60	7.70	9.40	0.25 ^a
Ox, ppb	321	42.98	8.99	38.40	43.40	48.70	47	46.74	8.99	41.50	46.20	53.60	0.015 ^a
SPM, µg/m ³	321	23.95	11.74	16.00	21.60	29.30	47	53.14	33.39	34.30	44.60	56.80	<0.01 ^a
Average Temperature, °C	321	14.78	4.4	11.5	14.9	18.6	47	17.4	3.9	15.4	18.0	20.3	<0.01 ^a
Maximum Temperature, °C	321	18.88	4.7	15.3	19.1	22.6	47	22.1	4.2	19.5	22.4	25.6	<0.01 ^a
Minimum Temperature, °C	321	11.13	4.7	7.5	10.8	15.0	47	13.1	4.2	10.5	13.9	16.6	<0.01 ^a
Humidity, %	321	67.49	13.3	58.0	68.0	76.0	47	67.2	8.4	62.0	68.0	74.0	0.56 ^b

SD, standard deviation; SPM, small particulate matter.

P value for ^aMann-Whitney U test and ^bt-test.

1.067–1.690), 1.318 (95% CI, 1.047–1.660), and 1.344 (95% CI, 1.071–1.687), respectively. The association was particularly prominent in school children, with ORs of 1.837 (95% CI, 1.212–2.786) for lag day 3 and 1.820 (95% CI, 1.179–2.808) for lag day 4. In preschool children, the crude ORs were around 1.1 for lag day 2 through lag day 5, without statistical significance.

As for respiratory diseases, AD events at lag day 0, lag day 1, and lag day 2 were significantly associated with emergency department visits by all children with ORs of 1.182 (95% CI, 1.084–1.288), 1.277 (95% CI, 1.169–1.393), and 1.262 (95% CI, 1.156–1.377), respectively. ORs for lag 01 and lag 02 were also increased. Unlike with bronchial asthma, however, the association was seemed stronger in preschool children than in school children, with ORs of 1.244 (95% CI, 1.128–1.373) for lag day 0, 1.314 (95% CI, 1.189–1.452) for lag day 1, and 1.273 (95% CI, 1.152–1.408) for lag day 2,

respectively. In schoolchildren, there was no statistically significant association except for lag day 2 (crude OR 1.227; 95% CI, 1.026–1.467).

We further adjusted for meteorological data and other air pollutants in our basic, single-pollutant, and multi-pollutant models. We also adjusted for spherical extinction coefficients in addition to the covariates in the basic model. These adjustments did not change the results substantially. In Table 3, we present the adjusted ORs of the association between AD days and emergency department visits due to bronchial asthma in school children, because a statistically significant association was observed in the crude analysis. Likewise, we present the adjusted ORs of the association between AD days and visits due to respiratory diseases in preschool children in Table 4. Although adjustment for covariates slightly attenuated the ORs, the results were again similar to those obtained using the basic model.




Figure 3. Crude associations between Asian dust and emergency department visits due to bronchial asthma (A) and respiratory diseases (B) by age group (vertical bar represents 95% confidence interval).

Table 3. Association between Asian dust and emergency department visits by bronchial asthma among school children in the basic, single-pollutant, and multi-pollutant models

	Basic model	Single-pollutant model (Basic model+single-pollutant) Adding to basic model, one air pollutant was adjusted			Multi-pollutant model (Single-pollutant model+other pollutant) Adding to basic model, two air pollutants were adjusted			Adding to basic model
	Adjusted by temperature+humidity OR (95% CI)	SO ₂ OR (95% CI)	NO ₂ OR (95% CI)	Ox OR (95% CI)	SO ₂ +NO ₂ OR (95% CI)	SO ₂ +Ox OR (95% CI)	NO ₂ +Ox OR (95% CI)	Spherical Extinction Coefficients OR (95% CI)
Lag 0	1.291 (0.850–1.961)	1.173 (0.760–1.811)	1.298 (0.854–1.974)	1.150 (0.738–1.790)	1.164 (0.754–1.798)	1.131 (0.725–1.765)	1.127 (0.724–1.757)	1.343 (0.867–2.081)
Lag 1	1.599 (1.032–2.478)	1.446 (0.911–2.294)	1.634 (1.048–2.546)	1.454 (0.916–2.307)	1.459 (0.918–2.316)	1.409 (0.883–2.250)	1.461 (0.921–2.317)	1.701 (1.080–2.681)
Lag 2	1.394 (0.891–2.182)	1.206 (0.740–1.964)	1.465 (0.915–2.344)	1.233 (0.765–1.990)	1.265 (0.770–2.080)	1.169 (0.714–1.914)	1.297 (0.798–2.110)	1.436 (0.898–2.298)
Lag 3	1.787 (1.174–2.721)	1.675 (1.088–2.579)	1.853 (1.205–2.848)	1.670 (1.080–2.581)	1.745 (1.127–2.701)	1.644 (1.061–2.547)	1.727 (1.113–2.680)	1.845 (1.204–2.826)
Lag 4	1.807 (1.163–2.809)	1.749 (1.121–2.730)	1.831 (1.175–2.852)	1.693 (1.076–2.666)	1.790 (1.145–2.799)	1.717 (1.088–2.710)	1.703 (1.082–2.680)	1.831 (1.173–2.857)
Lag 5	0.894 (0.550–1.453)	0.885 (0.542–1.442)	0.896 (0.551–1.458)	0.877 (0.538–1.430)	0.901 (0.552–1.469)	0.877 (0.538–1.432)	0.891 (0.547–1.451)	0.895 (0.550–1.457)
Lag 01	1.329 (0.915–1.930)	1.202 (0.809–1.786)	1.342 (0.922–1.955)	1.195 (0.800–1.787)	1.198 (0.805–1.781)	1.164 (0.775–1.748)	1.180 (0.789–1.765)	1.395 (0.947–2.056)
Lag 02	1.136 (0.794–1.625)	0.979 (0.659–1.454)	1.153 (0.800–1.663)	0.976 (0.655–1.455)	0.990 (0.666–1.471)	0.930 (0.617–1.401)	0.979 (0.656–1.459)	1.171 (0.804–1.705)
Lag 03	1.173 (0.840–1.639)	1.036 (0.718–1.485)	1.195 (0.848–1.684)	1.021 (0.699–1.491)	1.054 (0.730–1.523)	0.984 (0.668–1.450)	1.026 (0.702–1.500)	1.217 (0.858–1.726)
Lag 04	1.131 (0.822–1.557)	1.012 (0.716–1.431)	1.146 (0.828–1.587)	0.998 (0.699–1.423)	1.025 (0.725–1.449)	0.967 (0.672–1.389)	1.000 (0.701–1.426)	1.175 (0.844–1.635)
Lag 05	1.020 (0.739–1.408)	0.897 (0.631–1.276)	1.029 (0.741–1.429)	0.889 (0.623–1.269)	0.910 (0.640–1.295)	0.854 (0.592–1.231)	0.895 (0.627–1.278)	1.053 (0.756–1.467)

CI, confidence interval; OR, odds ratio.

We additionally performed a dose-response analysis of the basic model, classifying AD days using the AD extinction coefficients. In the case of bronchial asthma, the ORs were also statistically significant for lag day 3 or lag day 4 (eTable 1). For respiratory diseases, the ORs at lag day 0, lag day 1, and lag day 2 were elevated (eTable 2).

We also carried out a sensitivity analysis in which we excluded visits within 4 weeks of the initial visit. This gave us larger adjusted ORs than the previous results (eTable 3). We also compared whole-year data with those obtained between March and May; the ORs were lower with the whole-year data (eTable 4).

DISCUSSION

In this study, we found that exposure to AD led to increased emergency department visits by children. The association might vary between school and preschool children, though we could not detect a statistical interaction due to the small sample size. The association between AD exposure and increased emergency department visits due to bronchial asthma was might be stronger in school children than in preschool children, and the effect of exposure was manifested in a delayed fashion.

AD particles contain rock-forming minerals, such as quartz and feldspar, and argillites, such as mica, kaolinite, and green

Table 4. Association between Asian dust and emergency department visits by respiratory diseases among preschool children in the basic, single-pollutant, and multi-pollutant models

	Basic model	Single-pollutant model (Basic model+single pollutant) Adding to basic model, one air pollutant was adjusted			Multi-pollutant model (Single-pollutant model+other pollutant) Adding to basic model, two air pollutants were adjusted			Adding to basic model
	Adjusted by temperature+humidity OR (95% CI)	SO ₂ OR (95% CI)	NO ₂ OR (95% CI)	Ox OR (95% CI)	SO ₂ +NO ₂ OR (95% CI)	SO ₂ +Ox OR (95% CI)	NO ₂ +Ox OR (95% CI)	Spherical Extinction Coefficients OR (95% CI)
Lag 0	1.245 (1.129–1.374)	1.217 (1.099–1.347)	1.255 (1.137–1.385)	1.256 (1.129–1.396)	1.216 (1.099–1.347)	1.242 (1.117–1.382)	1.250 (1.124–1.390)	1.335 (1.202–1.483)
Lag 1	1.312 (1.186–1.451)	1.284 (1.156–1.426)	1.334 (1.204–1.477)	1.337 (1.198–1.492)	1.295 (1.165–1.439)	1.320 (1.182–1.475)	1.339 (1.200–1.495)	1.420 (1.276–1.580)
Lag 2	1.272 (1.148–1.410)	1.242 (1.117–1.382)	1.310 (1.178–1.456)	1.281 (1.148–1.429)	1.273 (1.143–1.418)	1.264 (1.132–1.411)	1.301 (1.165–1.453)	1.335 (1.201–1.484)
Lag 3	1.090 (0.980–1.213)	1.069 (0.960–1.190)	1.103 (0.990–1.229)	1.080 (0.969–1.204)	1.085 (0.973–1.209)	1.072 (0.961–1.195)	1.090 (0.977–1.216)	1.102 (0.990–1.227)
Lag 4	0.991 (0.891–1.102)	0.985 (0.885–1.095)	0.996 (0.895–1.108)	0.982 (0.882–1.094)	0.994 (0.893–1.106)	0.986 (0.885–1.098)	0.986 (0.886–1.098)	0.997 (0.896–1.109)
Lag 5	0.789 (0.708–0.880)	0.792 (0.710–0.884)	0.791 (0.710–0.883)	0.791 (0.709–0.882)	0.799 (0.716–0.892)	0.792 (0.710–0.884)	0.795 (0.713–0.887)	0.785 (0.704–0.875)
Lag 01	1.235 (1.132–1.348)	1.210 (1.103–1.327)	1.249 (1.144–1.364)	1.258 (1.142–1.386)	1.213 (1.106–1.330)	1.242 (1.126–1.369)	1.256 (1.140–1.383)	1.320 (1.203–1.448)
Lag 02	1.144 (1.054–1.241)	1.111 (1.018–1.214)	1.161 (1.068–1.262)	1.153 (1.050–1.265)	1.120 (1.025–1.223)	1.133 (1.030–1.245)	1.155 (1.052–1.267)	1.210 (1.109–1.321)
Lag 03	1.062 (0.983–1.147)	1.024 (0.943–1.112)	1.074 (0.993–1.163)	1.049 (0.962–1.145)	1.032 (0.950–1.122)	1.031 (0.943–1.126)	1.052 (0.964–1.148)	1.101 (1.015–1.194)
Lag 04	1.052 (0.978–1.132)	1.019 (0.943–1.102)	1.063 (0.987–1.145)	1.039 (0.959–1.127)	1.027 (0.950–1.110)	1.025 (0.944–1.112)	1.042 (0.961–1.130)	1.082 (1.002–1.168)
Lag 05	1.003 (0.934–1.078)	0.970 (0.899–1.046)	1.012 (0.940–1.089)	0.985 (0.912–1.064)	0.978 (0.906–1.055)	0.969 (0.896–1.049)	0.990 (0.916–1.070)	1.020 (0.947–1.098)

CI, confidence interval; OR, odds ratio.

mudstones. However, they also have sulfur oxidants, nitrogen oxidants, ammonium, and microorganisms attached to them, and these may have adverse health effects, including allergic reactions²⁷ and inflammation of the bronchi.²⁸ In Taiwan, Yu et al reported that the relative rate of clinic visits by children for respiratory diseases decreased significantly (by 0.74 to 0.99 times) in most districts during AD-storm periods, while it rose by 1.01 to 1.11 times in more than half of the districts during the period following AD storm, especially in school children.⁹ In contrast, Ueda et al reported no significant association between AD exposure and asthma hospitalization among children in Fukuoka, Japan.¹⁵

A possible explanation for our findings of a stronger association between AD exposure and emergency department visits due to bronchial asthma in school children than preschoolers is that the former participate in more outdoor activities and subsequently have greater exposure. Bronchial asthma is also more easily diagnosed in school children, and it is sometimes indistinguishable from other respiratory diseases in preschool children, such as bronchiolitis. Furthermore, preschool children with bronchial asthma might be under more careful management by their parents than school children.

Our study also revealed a delayed effect of AD exposure, with the strongest effect observed 3 or 4 days after exposure. This is consistent with some earlier findings.^{29,30} Yang et al reported that the effects of dust storms on admissions for asthma were most prominent 2 days after the event; however, this result was not statistically significant.²⁹ Meng et al reported that dust events were significantly associated with total respiratory hospitalizations 3 days later in both boys and girls.³⁰ This delayed effect may be due to physiological reasons; the respiratory function of school children is more mature than that of preschool children, so the older children can tolerate asthmatic symptoms to a greater extent. Another explanation may be delayed emergency department visits, as school children may not visit on the day when they notice

the symptoms because they do not want to miss school. Moreover, cough or wheeze and dyspnea with bronchial asthma are likely to worsen at night time.

As for the association between AD exposure and respiratory diseases, this was stronger in preschool children than in school children, and the effects were immediate. Although this association has been reported previously, the findings have been inconsistent in terms of susceptibility by age group and lag period. A study in Taipei showed that the rate of clinic visits for respiratory diseases during the weeks following AD events increased by 2.54% in preschool children and 5.03% in school children.⁸ In another study, post-AD days 1 through 4 saw significantly higher mean numbers of pneumonia admissions than non-AD days.¹¹ Yu et al reported a significantly increased risk of respiratory diseases in the week after AD, especially in school children. The largest percentage change in preschool children was at lag day 2 and lag day 3 (2.12% and 2.19%, respectively), and the largest change in school children was at lag day 3 (3.17%).¹⁰

The stronger association between AD exposure and respiratory diseases in preschool children than in school children observed in our study may be explained by the under-developed respiratory system in preschool children and difficulty in making differential diagnosis of wheezing illnesses, including bronchiolitis and other respiratory diseases, as well as bronchial asthma.

One of the strengths of the present study was the use of LIDAR to assess AD exposure. In Japan, AD days are generally assessed on the basis of visual observations by the Japan Meteorological Agency. These observations are done every hour, and an AD day is recorded when AD is observed for at least 1 hour on that day.³¹ In contrast, LIDAR measurements allow the quantitative evaluation of the level of AD particles as a dust extinction coefficient.¹⁸ However, no cutoff point has been established to distinguish between AD days and non-AD days. Kanatani et al used a 24-hour average dust extinction coefficient of 0.1/km or more to

define mild AD days.¹⁴ Ueda et al went further to define mild AD (dust extinction coefficient $\leq 0.066/\text{km}$), moderate AD ($0.066/\text{km}$ – $0.105/\text{km}$), and heavy AD days ($>0.105/\text{km}$).²⁶ Establishing a proper definition of AD days is warranted in the future.

The high regional representativeness of our study is another strength. The NMPEM Center is the only hospital in the study area that children can visit during the night, and approximately 90% of patients who access this center live in the covered area. However, we could not obtain patients' residential address because of laws governing the protection of personal information.

Time-stratified case-crossover analysis obviates the need to consider potential confounding by sex, age, and daily habits, which do not change over short periods, and also controls for the effects of long-term trends, seasonality, and the day of the week. However, a potential limitation of our study is that inflammation of the bronchi may continue for several days to several weeks after exposure, and inflammation makes the bronchi more vulnerable to environmental changes; therefore, those children with bronchial inflammation are more likely to revisit medical centers. To adjust for this, we repeated our analyses after excluding visits within 4 weeks of the initial visit and found that the results were substantially unchanged. In the sensitivity analysis, we compared the ORs in the whole year with those obtained between March and May, and found that the former was lower than the latter. Because most of the AD comes in the spring, we thought that the addition of seasons other than spring into the analysis would only attenuate the association between AD and outcomes.

Another potential limitation is a misclassification of exposure. LIDAR cannot measure the vertical distribution of AD any lower than 120 meters above the ground, so we cannot discount the possibility that the concentrations of particles our subjects actually inhaled were different from those measured. Nor did we obtain information on the patients' daily activities, so we could not take these into consideration. As for the horizontal distribution of AD, the LIDAR we used was located in Omura City, which is approximately 30 km from the center of Nagasaki City. However, we believe that any resulting misclassification would have been non-differential and would have made the observed association between AD exposure and outcomes weaker than the true association. Misclassification of the outcomes is also possible. Had the emergency physicians known that AD was on its way, it could have led to an information bias. However, if the elevated ORs were due only to such bias, the results would have been similar between preschool and school children, which was not the case. Another possible limitation we should mention is that, although the emergency medical center is open from 20:00 until 06:00, we did not obtain the exact times of patients' visits. This could also have led to outcome misclassification, but because any such misclassification was independent of

exposure, it would have been non-differential. In addition, when comparing our results with those from other countries, we have to consider the differences in diameter and the chemical composition of AD and in the attached metals.²⁸

Conclusions

Our study showed that exposure to AD was associated with increased emergency department visits due to bronchial asthma in school children and with increased visits due to respiratory diseases in preschool children.

ONLINE ONLY MATERIALS

eTable 1. Association between AD and emergency department visits due to bronchial asthma by categories of the AD extinction coefficients.

eTable 2. Association between AD and emergency department visits due to respiratory diseases by categories of the AD extinction coefficients.

eTable 3. Association between AD and emergency department visits by bronchial asthma among school children; all subjects vs subjects excluding those revisiting within 4 weeks.

eTable 4. Association between AD and emergency department visits by bronchial asthma among school children; comparison March–May vs whole year. Abstract in Japanese.

ACKNOWLEDGEMENTS

This work was conducted as part of a study on the health effects of AD conducted by Japan's Ministry of the Environment. The lead author (Takahiro Nakamura) received a grant from Dr. Takeshi Yanase of Toho University.

Conflicts of interest: None declared.

REFERENCES

1. Hashizume M, Ueda K, Nishiwaki Y, Michikawa T, Onozuka D. Health effects of AD events: a review of the literature. *Nihon Eiseigaku Zasshi*. 2010;65:413–21 (in Japanese).
2. Uno I, Eguchi K, Yumimoto K, Takemura T, Shimizu A, Uematsu M, et al. AD transported one full circuit around the globe. *Nat Geosci*. 2009;2:557–60.
3. Huebert BJ, Bates T, Russel PB, Shi G, Kim YJ, Kawamura K, et al. An overview of ACE-Asia: strategies for quantifying the relationships between Asian aerosols and their climatic impacts. *J Geophys Res*. 2003;108.
4. Sun J, Zhang M, Liu T. Spatial and temporal characteristics of dust storms in China and its surrounding regions, 1960–1999: relations to source area and climate. *J Geophys Res*. 2001;106:10325.
5. Vanos JK. Children's health and vulnerability in outdoor microclimates: a comprehensive review. *Environ Int*. 2015;76:1–15.

6. Darrow LA, Klein M, Flanders WD, Mulholland JA, Tolbert PE, Strickland MJ. Air pollution and acute respiratory infections among children 0–4 years of age: an 18-year time-series study. *Am J Epidemiol.* 2014;180:968–77.
7. Patel MM, Miller RL. Air pollution and childhood asthma: recent advances and future directions. *Curr Opin Pediatr.* 2009;21:235–42.
8. Chien LC, Yang CH, Yu HL. Estimated effects of AD storms on spatiotemporal distributions of clinic visits for respiratory diseases in Taipei children (Taiwan). *Environ Health Perspect.* 2012;120:1215–20.
9. Yu HL, Yang CH, Chien LC. Spatial vulnerability under extreme events: a case of AD storm's effects on children's respiratory health. *Environ Int.* 2013;54:35–44.
10. Yu HL, Chien LC, Yang CH. AD storm elevates children's respiratory health risks: a spatiotemporal analysis of children's clinic visits across Taipei (Taiwan). *PLoS One.* 2012;7:e41317.
11. Kang JH, Keller JJ, Chen CS, Lin HC. AD storm events are associated with an acute increase in pneumonia hospitalization. *Ann Epidemiol.* 2012;22:257–63.
12. Chan CC, Chuang KJ, Chen WJ, Chang WT, Lee CT, Peng CM. Increasing cardiopulmonary emergency department visits by long-range transported AD storms in Taiwan. *Environ Res.* 2008;106:393–400.
13. Wang CH, Chen CS, Lin CL. The threat of AD storms on asthma patients: a population-based study in Taiwan. *Glob Public Health.* 2014;9:1040–52.
14. Kanatani KT, Ito I, Al-Delaimy WK, Adachi Y, Mathews WC, Ramsdell JW. Desert dust exposure is associated with increased risk of asthma hospitalization in children. *Am J Respir Crit Care Med.* 2010;182:1475–81.
15. Ueda K, Nitta H, Odajima H. The effects of weather, air pollutants, and AD on hospitalization for asthma in Fukuoka. *Environ Health Prev Med.* 2010;15:350–7.
16. Kim SW, Yoon SC, Kim J, Kang JY, Sugimoto N. AD event observed in Seoul, Korea, during 29–31 May 2008: analysis of transport and vertical distribution of dust particles from lidar and surface measurements. *Sci Total Environ.* 2010;408:1707–18.
17. Hong CS, Lee KH, Kim YJ, Iwasaka Y. LIDAR measurements of the vertical aerosol profile and optical depth during the ACE-Asia 2001 IOP, at Gosan, Jeju Island, Korea. *Environ Monit Assess.* 2004;92:43–57.
18. Shimizu A, Sugimoto N, Matsui I, Arao K, Uno I, Murayama T, et al. Continuous observations of AD and other aerosols by polarization lidars in China and Japan during ACE-Asia. *J Geophys Res.* 2004;109.
19. Nagasaki Web City. Statistics data. 2015 National Census Information [cited 2015 July 1]. Available from: <http://www.city.nagasaki.lg.jp.e.jc.hp.transer.com/index.html>.
20. Nakamura T, Hashizume M, Ueda K, Kubo T, Shimizu A, Okamura T, et al. The relationship between AD events and out-of-hospital cardiac arrests in Japan. *J Epidemiol.* 2015;25:289–96.
21. Shimizu A, Sugimoto N, Matsui I, Mori I, Nishikawa M, Kido M. Relationship between lidar-derived dust extinction coefficients and mass concentrations in Japan. *Sola.* 2011;7A:1–5.
22. Yamazaki S, Shima M, Yoda Y, Oka K, Kurosaka F, Shimizu S, et al. Association of ambient air pollution and meteorological factors with primary care visits at night due to asthma attack. *Environ Health Prev Med.* 2013;18:401–6.
23. Maclure M, Mittleman MA. Should we use a case-crossover design? *Annu Rev Public Health.* 2000;21:193–221.
24. Maclure M. The case-crossover design: a method for studying transient effects on the risk of acute events. *Am J Epidemiol.* 1991;133:144–53.
25. Yuuhei HY, Motohiro E, Naomi K. Pediatric Guideline for the Treatment and Management of Asthma 2012. 2012:2–44 (in Japanese).
26. Ueda K, Shimizu A, Nitta H, Inoue K. Long-range transported AD and emergency ambulance dispatches. *Inhal Toxicol.* 2012;24:858–67.
27. Ichinose T, Yoshida S, Hiyoshi K, Sadakane K, Takano H, Nishikawa M, et al. The effects of microbial materials adhered to Asian sand dust on allergic lung inflammation. *Arch Environ Contam Toxicol.* 2008;55:348–57.
28. Mori I, Nishikawa M, Tanimura T, Quan H. Change in size distribution and chemical composition of kosa (AD) aerosol during long-range transport. *Atmos Environ.* 2003;37:4253–63.
29. Yang CY, Tsai SS, Chang CC, Ho SC. Effects of AD storm events on daily admissions for asthma in Taipei, Taiwan. *Inhal Toxicol.* 2005;17:817–21.
30. Meng Z, Lu B. Dust events as a risk factor for daily hospitalization for respiratory and cardiovascular diseases in Minqin, China. *Atmos Environ.* 2007;41:7048–58.
31. Ueda K, Shimizu A, Inoue K. Exposure assessment by light detection and ranging for AD aerosol in epidemiological studies to investigate its health effects. *Eurozoru Kenkyu.* 2014;29:7 (in Japanese).