

1 **A randomised phase III trial of skin closure by subcuticular suture versus skin stapling for**  
2 **prevention of incisional surgical site infection after elective colorectal cancer surgery**

3  
4  
5 Shin Kobayashi, MD<sup>1,5,8</sup>, Masaaki Ito, PhD<sup>1</sup>, Seichiro Yamamoto, PhD<sup>2</sup>, Yusuke Kinugasa, PhD<sup>3</sup>,  
6 Masanori Kotake, MD<sup>4</sup>, Yoshihisa Saida, PhD<sup>5</sup>, Takaya Kobatake, PhD<sup>6</sup>, Takeharu Yamanaka, PhD<sup>7</sup>,  
7 Norio Saito, PhD<sup>1</sup>, Yoshihiro Moriya, PhD<sup>2</sup>

8  
9 <sup>1</sup>Department of Colorectal Surgery, National Cancer Center Hospital East

10 <sup>2</sup>Department of Colorectal Surgery, National Cancer Center Hospital

11 <sup>3</sup>Department of Colon and Rectal Surgery, Shizuoka Cancer Center

12 <sup>4</sup>Department of Gastroenterological Surgery, Ishikawa Prefectural Central Hospital

13 <sup>5</sup>Department of Surgery, Toho University Ohashi Medical Center

14 <sup>6</sup>Department of Gastroenterological Surgery, National Hospital Organization Shikoku Cancer Center

15 <sup>7</sup>Department of Biostatistics, Yokohama City University

16 <sup>8</sup>Department of Surgery, Tokyo Metropolitan Komagome Hospital

17  
18 Corresponding author: Masaaki Ito

19 6-5-1 Kashiwanoha, Kashiwa, Chiba 277-8577, Japan

20 Tel: +81-4-7133-1111; Fax: +81-4-7131-9960

21 E-mail: maito@east.ncc.go.jp

22  
23 Funding: A Grant-in-Aid for Cancer Research in Japan.

24  
25 Category of manuscript: Randomized clinical trial

26  
27 This paper does not have previous communication to a society or meeting.

1 **Abstract**

2

3 **Background:** Whether subcuticular suture prevents incisional surgical site infection (SSI) after  
4 clean-contaminated surgery has not been clarified.

5

6 **Method:** Patients undergoing elective colorectal cancer surgery at one of 16 centers were randomised  
7 to receive either subcuticular sutures or skin stapling during the skin closure part of the procedure.

8 The occurrence of incisional SSI was the endpoint of interest. Also accounted for were the time  
9 required for wound closure, incidence of wound problems, postoperative length of stay, wound  
10 aesthetics, and patient satisfaction.

11

12 **Results:** A total of 1264 patients were enrolled in the study. The cumulative incidence of incisional  
13 SSI by day 30 after surgery did not differ significantly between the subcuticular suture group and the  
14 staple closure group (8.7% vs 9.8%, respectively;  $P=0.58$  by chi-square test) (odds ratio: 0.88, 95% CI  
15 0.60-1.29). However, a comparison of cumulative incidence curves revealed a significantly longer  
16 time to SSI in the subcuticular suture group ( $P=0.019$  by log-rank test) (hazard ratio: 0.66, 95% CI  
17 0.45-0.97). Wound problems ( $P=0.48$ ), wound aesthetics ( $P=0.18$ ), and postoperative length of stay  
18 ( $P=0.51$ ) did not differ significantly between groups, and the time required for wound closure was 5  
19 minutes longer in the subcuticular suture group ( $P<0.001$ ). However, patients in the subcuticular  
20 suture group were significantly more satisfied with their wound (52% vs 43%,  $P=0.002$ ).

21

22 **Conclusion:** Subcuticular suture does not seem to significantly reduce the risk of incisional SSI  
23 occurring within 30 days after surgery. Registration number: UMIN000004001.

24

25

## 1 **Introduction**

2 Surgical site infection (SSI) is one of the most common postoperative complications encountered,  
3 with a reported incidence of 500,000 cases per year in the United States.<sup>1</sup> SSIs have been reported to  
4 account for 15% of all nosocomial infections and up to 38% of such infections in surgical patients.<sup>2</sup>  
5 Furthermore, SSIs have been reported to account for 7.3 extra days of hospital stay and for \$3,152 in  
6 additional costs.<sup>3</sup> Most importantly, 77% of postoperative mortalities have been related to SSIs.<sup>4</sup>  
7 Therefore, preventing SSIs is crucial to safeguarding patients and containing medical costs. Incisional  
8 SSIs in particular account for two thirds of all SSIs.

9

10 Subcuticular suture is a method of wound closure by which the opposing dermal layers at each edge  
11 are attached with absorbable sutures buried beneath the skin surface (figure 1a). This method was first  
12 described by Halsted in 1890 and has been the preferred method for wound closure in clean surgery.<sup>5</sup>  
13 Because no foreign material reaches beyond the epidermis, this method of closure causes no undue  
14 skin inflammation, obviates the need for postoperative removal of the suture material, and enhances  
15 wound aesthetics.<sup>6-8</sup> The reported rate of incisional SSI is lower with subcuticular suture closure than  
16 with staple closure.<sup>6,9,10</sup>

17

18 The prevention of incisional SSIs by subcuticular suture was very recently tested in the context of  
19 gastroenterological surgery,<sup>11</sup> and the incidence of incisional SSIs appeared to be statistically reduced  
20 in cases of colorectal surgery in particular. However, the sample group was relatively small, and the  
21 procedures were all open surgeries. The study result pointed to the need for a large-scale randomised  
22 controlled trial that would include both open and laparoscopic surgeries to reflect current surgical  
23 practice. Thus, the present multi-centre study was conducted to examine whether subcuticular suture,  
24 in comparison to metal staple closure, reduces the incidence of incisional SSI after surgery for  
25 colorectal cancer.

26

27

## 1 **Methods**

### 2 **Study design**

3 Eligible patients 1) were scheduled for elective colorectal cancer surgery with or without ileostomy or  
4 colostomy; 2) had an Eastern Co-operative Oncology Group performance status (ECOG PS) of 0 or 1;  
5 and 3) showed adequate organ function according to the following laboratory data: white blood cell  
6 count  $> 3,000$  and  $< 10,000/\mu\text{L}$ ; neutrophil count  $\geq 1,500/\mu\text{L}$ ; hemoglobin  $\geq 9.0$  g/dL; platelet count  $\geq$   
7  $70,000/\mu\text{L}$ ; total bilirubin  $\leq 1.5$  mg/dL; AST  $\leq 100$  IU/L; ALT  $\leq 100$  IU/L; serum creatinine  $\leq 1.5$   
8 mg/dL; and serum albumin  $\geq 2.8$  g/dL; and 4) provided written informed consent.

9

10 Eligible patients were assigned randomly to skin closure by subcuticular suture or by skin stapling.  
11 The primary endpoint was the occurrence of postoperative incisional SSI, and the cumulative  
12 incidence of such SSI was determined. Other factors investigated were 1) wound problems, especially  
13 excessive exudation or dehiscence; 2) length of postoperative hospital stay; 3) time required to close  
14 the wound; 4) wound aesthetics; and 5) patient satisfaction. This study was approved by the  
15 institutional review board (IRB) of each center involved and registered in the UMIN Clinical Trials  
16 Registry (UMIN000004001).

17

### 18 **Randomisation and masking**

19 Patients were recruited by participating surgeons and enrolled preoperatively. Randomisation was  
20 done at a 1:1 allocation ratio to balance treatment over the following factors: institution, type of  
21 surgery (open vs. laparoscopic), and location of the tumour (colon vs. rectum). Surgeons were notified  
22 of the allocation by telephone before surgery; neither the patient nor the investigators were blinded to  
23 the allocation. These processes were managed by the data center located at National Cancer Center  
24 Hospital East.

25

### 26 **Surgical procedure**

27 After fascial closure, the incision was irrigated with the appropriate amount of warm saline,  
28 approximately 500 mL for open surgery and 100 mL for laparoscopic surgery. The incision was then

1 closed with either subcuticular sutures or skin staples. When subcuticular sutures were used, the  
2 dermal layers of each edge were properly attached to each other with suture strings buried beneath the  
3 skin surface (figure 1a); 4-0 or 5-0 absorbable mono-filament suture material was used. When skin  
4 staples were used, the dermis of each edge was properly attached at intervals of 10-15 mm (figure 1b).  
5 For standardisation, the participating surgeons were requested to learn the procedures described above  
6 through lectures and by viewing an instructional video provided by the authors before the start of the  
7 study.

8

### 9 **Perioperative care**

10 Because the study was a multi-institutional trial, perioperative care was not exactly the same from  
11 centre to centre. The institutions were surveyed before the trial by means of a questionnaire, and it  
12 was confirmed that each institution largely follows the Centers for Disease Control and Prevention  
13 (CDC) guideline for perioperative care.<sup>4</sup> The approach to standard perioperative care as described in  
14 the CDC guideline was also provided to each institution at conferences and in the study protocol.

15

### 16 **SSI surveillance**

17 Patients' surgical wounds were inspected daily during the period of hospitalisation. Incisional SSI was  
18 suspected if there was any purulent discharge, pain or tenderness, localised swelling, redness, or heat.  
19 The fluid or tissue was aseptically obtained and cultured, and if infectious organisms were isolated,  
20 SSI was diagnosed, and the superficial incision was opened by the surgeon.<sup>4</sup> An SSI was identified by  
21 a surgeon other than the patient's primary surgery or by the institution's infection control team. Each  
22 participating surgeon attended lectures, meetings, and discussions to learn to identify SSIs on the  
23 basis of the CDC criteria. These sessions were held to standardise the surgeons' interpretation of the  
24 SSI definition as well as the surveillance at each institution. Upon hospital discharge, each patient was  
25 asked to respond to a questionnaire regarding his or her wound, and the incision was re-inspected at  
26 the outpatient clinic approximately 30 days after surgery. Therefore, the possibility of post-discharge  
27 SSI was tracked both by questionnaire and out-patient examination. The total number of incisional  
28 SSIs detected overall within 30 days after surgery, both during and after hospitalisation, was

1 determined.

2

### 3 **Evaluation of other factors**

4 Wound problems, particularly exudation and dehiscence, were surveyed in the same way as SSI. The  
5 total number of patients with excessive exudate or dehiscence within 30 days after surgery, both  
6 during and after hospitalisation, was recorded. Each patient's wound was inspected for aesthetics in  
7 the outpatient clinic 30 days after surgery by the nurse and surgeon and judged in accordance with a  
8 validated scar evaluation scale ranging from 1 (least aesthetic) to 6 (most aesthetic).<sup>12</sup> Patient  
9 satisfaction was self-reported on a scale ranging from 1 (least possible satisfaction) to 5 (greatest  
10 possible satisfaction).<sup>13</sup>

11

### 12 **Statistical analysis**

13 The study was first designed to detect a 6.0% reduction in the incidence of SSI from 14.0% in the  
14 staple group to 8% in the subcuticular suture group, with a two-sided significance level of 0.10 and  
15 power of 80%. Accordingly, the initial target sample size was 800 patients. However, upon  
16 consultation with a biostatistician (T.Y.), it was recommended that we change the two-sided  
17 significance level to 0.05 and the power to 90% to obtain a definitive result. This change necessitated  
18 enrollment of 1206 patients. We thus planned a sample size of 1240 patients. Between-group  
19 difference in the incidence of incisional SSI was analysed by chi-square test. Differences between  
20 numerical variables and binary variables were analysed by Mann-Whitney U-test and chi-square test,  
21 respectively. Between-group difference in the incidence of incisional SSI over time was analysed by  
22 log-rank test. All P values are two-tailed, and  $P < 0.05$  was considered statistically significant.  
23 Statistical analyses were performed with SAS for Windows ver. 9.3 and SPSS 11.0 J for Windows.

24

## 1 **Results**

2 In total, 1264 patients were enrolled from 16 centers in Japan between August 2010 and April 2012.  
3 Of these patients, 629 were allocated to abdominal closure by stapling and 635 to closure by  
4 subcuticular suture. Thirty-two patients, because of re-operation (staple group, n=16; subcuticular  
5 suture group, n=8), major protocol violation (subcuticular suture group, n=3), cancelled surgery  
6 (staple group, n=1; subcuticular suture group, n=2), withdrawn consent (subcuticular suture group,  
7 n=1), or ineligibility (subcuticular suture group, n=1) were excluded from the analysis (figure 2). The  
8 ineligibility was due to anemia (hemoglobin < 9.0 g/dL). Thus, the full analysis set comprised 612  
9 patients in the staple group and 620 in the subcuticular suture group. Baseline demographics and  
10 operative and postoperative characteristics of patients were well-balanced between the two groups  
11 (Table 1).

12

13 The study results are shown in Table 2. The incidence of incisional SSI at 30 days after surgery was  
14 9.8% in the staple group and 8.7% in the subcuticular suture group. This difference was not  
15 statistically significant (P=0.58). However, a significant between-group difference was found in the  
16 time to occurrence of SSI (figure 3; P=0.019), with a hazard ratio of 0.659 (95% CI, 0.449-0.968).  
17 The incidence of wound problems (6.9% vs. 5.8%; P=0.48), the postoperative length of stay (9 days  
18 vs. 9 days; P=0.51), and wound aesthetics (proportion of patients with highest possible score: 78% vs.  
19 80%; P=0.18) did not differ significantly between the two groups, but the time required to close the  
20 wound was significantly longer in the subcuticular suture group than in the staple group (1 minute vs.  
21 6.5 minutes; P<0.001). Regarding patient satisfaction, the proportion of patients giving the highest  
22 possible score was greater in the subcuticular suture group than in the staple group (P=0.002). In our  
23 subset analysis, subcuticular suture tended to be favored, but the difference was not significant (figure  
24 4).

## 1 **Discussion**

2 Subcuticular suture is a skin closure method that involves properly attaching opposite dermal layers  
3 with absorbable sutures buried beneath the skin surface. It is widely accepted as the procedure of  
4 choice in clean surgery because it is good for simple postoperative care, wound aesthetics, and wound  
5 scar prevention, and there is no need for removal of the stitches.<sup>5-7</sup> Subcuticular suture has been  
6 shown to be significantly better than stapling for preventing wound problems including incisional SSI  
7 after cardiovascular and orthopaedic surgery.<sup>6,9,14</sup> A group of gastrointestinal surgeons in Japan  
8 recently found and reported no difference between subcuticular suture and stapling in the incidence of  
9 wound problems after open gastrointestinal surgery.<sup>11</sup> However, a subcuticular suture-based decrease  
10 was found in the incidence of incisional SSI after open colorectal surgery in particular, although the  
11 number of colorectal surgeries was relatively low (n=277) and the procedure was limited to open  
12 surgery. We designed a large randomised controlled trial that would include both open and  
13 laparoscopic colorectal surgery to clarify whether subcuticular suture is of clinical benefit for  
14 preventing incisional SSI in cases of colorectal surgery. The present study was conducted in the  
15 context of colorectal surgery because the incidence of incisional SSI is higher in this context than in  
16 others.

17  
18 Physiologically, wound healing involves re-epithelialisation that is completed within 24-48 hours after  
19 skin closure, and dermal reconstruction takes place through angiogenesis and fibrogenesis within 3-4  
20 days.<sup>15-17</sup> We believe that subcuticular suture promotes attachment of each dermal layer, supports  
21 dermal reconstruction, and subsequently enhances wound healing. This series of proper wound  
22 healing events prevents tissue hypoxia and dead space formation, which encourage infection; thus, the  
23 surgical incision is protected from contamination. Therefore, we predicted subcuticular suture to be  
24 effective in preventing incisional SSI in colorectal surgery.

25  
26 Among our patients, the incidence of incisional SSI after subcuticular suture as of postoperative day  
27 30 did not differ significantly from that after stapling (8.7% vs. 9.8%, P=0.58 by chi-square test, odds  
28 ratio: 0.88, 95% CI 0.60-1.29). Subset analysis also indicated that subcuticular suture was not superior



1 over stapling for preventing SSI (figure 4). However, the cumulative risk of incisional SSI within 30  
2 days was shown to be significantly lower in the subcuticular suture group than in the staple group  
3 (figure 3) ( $P=0.019$  by log-rank test, hazard ratio: 0.66, 95% CI 0.45-0.97). These data indicate that  
4 subcuticular suture prevents incisional SSI to some extent but not unequivocally within the first 30  
5 days after surgery. Because the cumulative risk was shown to be lower in the subcuticular suture  
6 group, it is possible that the incidence of incisional SSI in the subcuticular suture group would also  
7 have been lower if the incisions had been judged beyond postoperative day 30. In the very least, it is  
8 evident that subcuticular suture is a feasible method of wound closure not only after clean surgery but  
9 also after colorectal surgery, for which the incidence of incisional SSI is relatively high.

10  
11 Wound problems including exudation and dehiscence, wound aesthetics, and postoperative hospital  
12 stay did not differ significantly between the two methods (Table 2). However, patient satisfaction,  
13 judged by the percentage of patients giving the highest possible score, was significantly greater in the  
14 subcuticular suture group than in the staple group ( $P=0.002$ ). Although wound aesthetics did not differ  
15 between the two groups, patients were more satisfied with subcuticular suture, probably because it  
16 does not require removal of the stitches or because of its natural appearance without foreign objects  
17 like staples. The longer time required to close the wounds by subcuticular suture was not unexpected,  
18 but the difference was only 5 minutes in an operation that was longer than 200 minutes. Given the  
19 context, this difference is minor.

20  
21 Tsujinaka and colleagues conducted the first randomised controlled trial investigating the influence of  
22 subcuticular suture on wound complications in gastrointestinal surgery.<sup>11</sup> Subcuticular suture was  
23 shown to decrease the incidence of incisional SSI in colorectal surgery in particular. However, the trial  
24 included only open surgeries, wound closure allocations were not well-balanced ( $n=176$  vs. 101), and  
25 tumour location was not considered among the allocation factors even though it is a significant  
26 prognostic factor for incisional SSI.<sup>18</sup> In comparison, the present study focused on colorectal cancer,  
27 involved both open and laparoscopic surgery, and balanced the patient allocation in terms of tumour  
28 location and type of surgery, both of which are significant factors related to the risk of incisional

1 SSI.<sup>18,19</sup> The proportion of laparoscopic surgeries was high in the present study, which was reflective  
2 of current surgical practice. Therefore, we believe that the study described herein clearly elucidates  
3 the influence of subcuticular suture on SSI following colorectal surgery. We believe this study to be  
4 complementary to the trial conducted by Tsujinaka and colleagues, which dealt only with open surgery.  
5 Together, the two studies show clearly that subcuticular suture is equivalent to skin stapling in terms  
6 of the risk of incisional SSI after gastrointestinal surgery.

7  
8 One of the limitations we face in discovering SSIs is post-discharge surveillance. Delgado-Rodriguez  
9 et al. reported development of SSIs in as many as 45.6% (103/226) of patients after discharge.<sup>20</sup>

10 However, in some centres, post-discharge SSI surveillance is based solely on a review of medical  
11 records or on a one-time medical examination in the outpatient clinic.<sup>6-10,14,21</sup> With the trend towards  
12 shorter hospital stays, some infections may be missed unless special attention is paid to post-discharge  
13 surveillance. However, in the present study, post-discharge SSI surveillance was achieved by means  
14 of patient questionnaire, outpatient medical examination, and review of the patient's medical records,  
15 which increased the reliability of the results. Another limitation was that the response to the patient  
16 questionnaire was 81.6%. However, this rate was fairly high in comparison to rates obtained in other  
17 surgical randomised controlled trials,<sup>22-24</sup> and it did not differ significantly between our two procedure  
18 groups (P=0.44). Therefore, we believe that this limitation did not substantially affect the  
19 interpretation of our data. The other limitation was that perioperative factors that affect the incidence  
20 of SSI were not exactly the same among the participating institutions. However, each institution was  
21 involved in several joint meetings to standardise the subcuticular suture technique, the perioperative  
22 care, and the SSI surveillance. Furthermore, institution was used as a stratification factor before  
23 randomisation. Therefore, the effects of inter-institutional inconsistencies were minimised.

24  
25 In conclusion, we found that the incidence of incisional SSI in patients treated by elective colorectal  
26 surgery does not differ significantly between closure of the skin by subcuticular suturing and closure  
27 by stapling. However, patients are more satisfied with wounds closed by subcuticular suture than with  
28 those closed with staples. Considering the increased patient satisfaction, we recommend subcuticular

1 suturing as the procedure of choice for skin closure after gastroenterological surgery.

## 1 **Acknowledgments**

2 This work was supported by a Grant-in-Aid for Cancer Research from the Ministry of Health, Labor  
3 and Welfare of Japan.

4 The authors would like to express their deepest appreciation for the discussions and suggestions given  
5 by Dr. Masanori Sugito, Akihiro Kobayashi, Yusuke Nishizawa (National Cancer Center Hospital  
6 East), Dr. Shin Fujita (National Cancer Center Hospital), Dr. Shinya Kusachi, Toshiyuki Enomoto  
7 (Toho University Ohashi Medical Center), Dr. Hiroyuki Bando (Ishikawa Prefectural Central  
8 Hospital), Dr. Masayuki Ohue, Shingo Noura, Tatsushi Shingai (Osaka Medical Center for Cancer and  
9 Cardiovascular Diseases), Dr. Akio Shiomi, Tomohiro Yamaguchi (Shizuoka Cancer Center), Dr.  
10 Yoshiro Kubo (Shikoku Cancer Center), Dr. Koji Komori, Yukihide Kanemitsu, Kenya Kimura,  
11 Takashi Kinoshita (Aichi Cancer Center), Dr. Kouki Ootsuka (Iwate Medical University), Dr.  
12 Toshihiko Sato (Yamagata Prefectural Central Hospital) , Dr. Keiichi Takahashi (Tokyo Metropolitan  
13 Komagome Hospital), Dr. Tomoharu Shimizu (Shiga University of Medical Science Hospital), Dr.  
14 Kazuhiko Yoshimatsu (Tokyo Women`s Medical University Medical Center East), Dr. Masayoshi  
15 Nakanishi, Yasutoshi Murayama (Kyoto Prefectural Medical University), Dr. Madoka Hamada  
16 (Kansai Medical University Hirakata Hospital), and Dr. Yasumasa Takii (Niigata Cancer Center  
17 Hospital). Also, the authors would like to express their deepest gratitude to Yuko Tanabe, Mieko  
18 Noguchi, Tomomi Murakoshi, Yuki Takahashi, and Noriko Okada for data management.

19

20

## 1 **References**

- 2 1. Nichols RL. Preventing surgical site infections: a surgeon's perspective. *Emerg Infect Dis* 2001; 7:  
3 220-24.
- 4 2. Emori TG, Gaynes RP. An overview of nosocomial infections, including the role of the  
5 microbiology laboratory. *Clin Microbiol Rev* 1993; 6: 428-42.
- 6 3. Martone WJ, Jarvis WR, Culver DH, Haley RW. Incidence and nature of endemic and epidemic  
7 nosocomial infections. In: Bennett JV, Brachman PS, eds. *Hospital Infections*. 3<sup>rd</sup> ed. Boston: Little,  
8 Brown and Co, 1992; 577-96.
- 9 4. Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR. Guideline for prevention of surgical  
10 site infection, 1999. *Infect Control Hosp Epidemiol* 1999; 20: 247-78.
- 11 5. Halsted WH. Radical cure of hernia. *Johns Hopkins Hospital Medical Journal* 1890; 1: 13.
- 12 6. Angelini GD, Butchart EG, Armistead SH, Breckenridge IM. Comparative study of leg wound skin  
13 closure in coronary artery bypass graft operations. *Thorax* 1984; 39: 942-45.
- 14 7. Lubowski D, Hunt D. Abdominal wound closure comparing the proximate stapler with sutures.  
15 *Aust N Z J Surg* 1985; 55: 405-6.
- 16 8. Ranaboldo CJ, Rowe-Jones DC. Closure of laparotomy wounds: skin staples versus sutures. *Br J*  
17 *Surg* 1992; 79: 1172-3.
- 18 9. Johnson RG, Cohn WE, Thurer RL, McCarthy JR, Sirois CA, Weintraub RM. Cutaneous closure  
19 after cardiac operations. *Ann Surg* 1997; 226: 606-12.
- 20 10. Johnson A, Young D, Reilly J. Caesarean section surgical site infection surveillance. *J Hosp Infect*  
21 2006; 64: 30-35.
- 22 11. Tsujinaka T, Yamamoto K, Fujita J, Endo S, Kawada J, Nakahira S. Subcuticular sutures versus  
23 staples for skin closure after open gastrointestinal surgery: a phase 3, multicenter, open-label,  
24 randomized controlled trial. *Lancet* 2013; 382: 1105-12.
- 25 12. Singer AJ, Arora B, Dagum A, Valentine S, Hollander JE. Development and validation of a novel  
26 scar evaluation scale. *Plast Reconstr Surg* 2007; 120: 1892-1897.
- 27 13. Hendriks AA, Vrieling MR, Smets EM, van Es SQ, De Haes JC. Improving the assessment of  
28 (in)patients' satisfaction with hospital care. *Med Care* 2001; 39: 270-83.
- 29 14. Shetty AA, Kumar VS, Morgan-Hough C, Georgeu GA, James KD, Nicholl JE. Comparing

- 1 wound complication rates following closure of hip wounds with metallic skin staples or subcuticular  
2 vicryl suture: a prospective randomized trial. *J Ortho Surg* 2004; 12: 191-93.
- 3 15. Lawrence WT. Wound healing biology and its application to wound management. In: O'Leary JP,  
4 ed. *The physiologic basis of surgery*, 3rd ed. Philadelphia: Lippincott Williams & Wilkins, 2002:  
5 107-32.
- 6 16. Li J, Chen J, Kirsner R. Pathophysiology of acute wound healing. *Clin Dermatol* 2007; 25: 9-18.
- 7 17. Singer AJ, Clark RAF. Cutaneous wound healing. *N Engl J Med* 1999; 341: 738-46.
- 8 18. Konishi T, Watanabe T, Kishimoto J, Nagawa H. Elective colon and rectal surgery differ in risk  
9 factors for wound infection: results of prospective surveillance. *Ann Surg* 2006; 244: 758-63.
- 10 19. Poon JT, Law WL, Wong IW, Fan JK, Lo OS. Impact of laparoscopic colorectal resection on  
11 surgical site infection. *Ann Surg* 2009; 249: 77-81.
- 12 20. Delgado-Rodriguez M, Gomez-Ortega A, Sillero-Arenas M, Liorca J. Epidemiology of  
13 surgical-site infections diagnosed after hospital discharge: a prospective cohort study. *Infect Control*  
14 *Hosp Epidemiol* 2001; 22: 24-30.
- 15 21. Fujita S, Saito N, Yamada T, Takii Y, Kondo K, Ohue M, et al. Randomized, multicenter trial of  
16 antibiotic prophylaxis in elective colorectal surgery. *Arch Surg* 2007; 142: 657-61.
- 17 22. Galmiche J, Hatlebakk J, Attwood S, Eil C, Fiocca R, Eklund S, et al. Laparoscopic antireflux  
18 surgery vs esomeprazole treatment for chronic GERD. The LOTUS randomized clinical trial. *JAMA*  
19 2011; 305: 1969-77.
- 20 23. Grant AM, Cotton SC, Boachie C, Ramsay CR, Krukowski ZH, Heading RC, et al. Minimal  
21 access surgery compared with medical management for gastro-oesophageal reflux disease: five year  
22 follow-up of a randomised controlled trial (REFLUX). *BMJ* 2013; 346: f1908.
- 23 24. Kornblith AB, Lan L, Archer L, Partridge A, Kimmick G, Hudis C, et al. Quality of life of older  
24 patients with early-stage breast cancer receiving adjuvant chemotherapy: a companion study to cancer  
25 and leukemia group B 49907. *J Clin Oncol* 2011; 29: 1022-8.
- 26 25. Shuster M. Comparing skin staples to sutures in an emergency department. *Can Fam Physician*  
27 1989; 35: 505-9.
- 28 26. Obermair A, Crandon A, Perrin L, Walsh T, Carrazo M, Nicklin J. Randomized trial of skin  
29 closure after laparotomy for gynaecological surgery. *ANZ J Surg.* 2007; 77: 460-3.

- 1 27. Clayer M, Southwood RT. Comparative study of skin closure in hip surgery. ANZ J Surg. 1991;
- 2 61: 363-5.
- 3

1 **Figure legends**

2

3 **Figure 1: Schematic drawings of skin closure**

4 **1-a) Subcuticular suturing**

5 The opposing skin layers are properly attached to each other by subcuticular suture.

6 **1-b) Skin stapling**

7 The opposing skin layers are attached to each other with skin staples.

8

9 **Figure 2: Study enrollment and randomisation**

10 A total of 1,264 patients were enrolled, 1,232 of whom were included in the final analysis.

11

12 **Figure 3: Cumulative risk of incisional SSI within 30 days after surgery**

13


14 **Figure 4: Subset analysis of the relative risk of incisional SSI**

15 Note that the relative risk of incisional SSI did not differ significantly between any of the subgroups.

16




1 Figure 1a



2

3 Figure 1b



4

5

1

2

Figure 2 Study enrolment and randomisation






Figure 3 Cumulative risk of incisional SSI within 30 days after surgery



No. at risk	3	6	9	12	15	18	21	24	27	30
Staples	612	611	590	571	563	558	554	554	553	553
Subcuticular suture	620	620	604	589	583	577	573	572	570	568

**Figure 4 Subgroup analysis**



**Table 1. Patient characteristics per study group**

<b>Characteristic</b>		<b>Staples (n=612)</b>		<b>Subcuticular suture (n=620)</b>	
Age (years)	median (range)	67	(25-91)	65	(30-91)
Gender (male)	n, (%)	335	(55%)	335	(54%)
BMI (kg/m <sup>2</sup> )	median (range)	22.6	(14.3- 38.2)	22.3	(14.6-34.3)
PS (0)	n, (%)	569	(93%)	577	(93%)
ASA score					
1, 2	n, (%)	595	(97%)	600	(97%)
3	n, (%)	17	(3%)	20	(3%)
DM	n, (%)	47	(8%)	64	(10%)
History of smoking	n, (%)	80	(13%)	66	(11%)
Type of surgery					
Open	n, (%)	121	(20%)	121	(20%)
Laparoscopic	n, (%)	491	(80%)	499	(80%)
Tumour location					
Colon	n, (%)	462	(76%)	457	(74%)
Rectum	n, (%)	150	(24%)	163	(26%)
Lowest body temperature during surgery (°C)	median (range)	35.8	(31.9-37.2)	35.9	(31.9-37.7)
Surgical drains	n, (%)	467	(76%)	469	(76%)
Prophylactic cephalosporin	n, (%)	609	(99%)	619	(99%)
Depth of subcutaneous fat tissue (cm)	median (range)	1.8	(0-5)	1.8	(0-5)
Operation time (min)	median (range)	206	( 60-690)	213	(81-621)
Blood loss (mL)	median (range)	29	(0-4700)	25	(0-3575)

BMI: body mass index, PS: performance status, ASA: American society of Anesthesiology, DM: diabetes mellitus

**Table 2 Postoperative outcomes according to the type of skin closure**

<b>Outcome</b>		<b>Staples (n=612)</b>		<b>Subcuticular suture (n=620)</b>		<b>P value</b>
Incisional SSI	n, (%)	60	(9.8%)	54	(8.7%)	0.58
Wound problem	n, (%)	42	(6.9%)	36	(5.8%)	0.48
Postoperative hospital stay (days)	median (range)	9	(4-77)	9	(5-76)	0.51
Wound aesthetics	n, (%) of patients with highest score	475	(78%)	500	(80%)	0.18
Patient satisfaction*	n, (%) of patients with highest score	211	(43%)	268	(52%)	0.002
Time to close wound (min)	median (range)	1	(0.5-25)	6.5	(0.5-30)	< 0.001

SSI: surgical site infection

\*total n=1005