

東邦大学学術リポジトリ

Toho University Academic Repository

タイトル	Pancreatic Insulinoma Associated with Stenosis of the Celiac Axis Due to Median Arcuate Ligament Compression
作成者（著者）	Kanazawa, Ken / Yoshino, Hiroshi / Kakumae Kojimahara, Yuki / Yoshikawa, Fukumi / Ando, Yasuyo / Kumashiro, Naoki / Uchino, Hiroshi / Hirose, Takahisa
公開者	The Medical Society of Toho University
発行日	2019.03.01
ISSN	21891990
掲載情報	Toho Journal of Medicine. 5(1). p.20 23.
資料種別	学術雑誌論文
内容記述	Case Report
著者版フラグ	publisher
JaLCDOI	info:doi/10.14994/tohojmed.2018 012
メタデータのURL	https://mylibrary.toho u.ac.jp/webopac/TD53416997

Case Report

Pancreatic Insulinoma Associated with Stenosis of the Celiac Axis Due to Median Arcuate Ligament Compression

Ken Kanazawa Hiroshi Yoshino* Yuki Kakumae-Kojimahara
Fukumi Yoshikawa Yasuyo Ando Naoki Kumashiro
Hiroshi Uchino and Takahisa Hirose

Department of Internal Medicine, Division of Diabetes, Metabolism and Endocrinology,
Toho University Graduate School of Medicine, Tokyo, Japan

ABSTRACT: Median arcuate ligament syndrome (MALS) is a rare stenotic vascular disorder caused by fibrous bands connecting the left and right crus of the diaphragm. A 42-year-old man was admitted to our hospital. Computed tomography revealed 20-mm mass at the tail of pancreas. He was diagnosed as having insulinoma. Angiography showed increased blood flow from dorsal pancreatic artery to insulinoma. Enhanced computed tomography revealed stenosis in the celiac artery. Thus, he was suspected to be suffering from MALS. MALS should be considered when angiography of area surrounding the pancreas reveals blood flow different from that observed in normal anatomy.

Toho J Med 5 (1): 20–23, 2019

KEYWORDS: median arcuate ligament syndrome, insulinoma

Median arcuate ligament syndrome (MALS) is a rare disorder caused by compression of celiac artery by the median arcuate ligament. MALS was first described in 1963.¹⁾ MALS is a rare vascular disorder caused by fibrous bands connecting the left and right crus of the diaphragm. Clinical symptoms are postprandial abdominal pain, vomiting, and weight loss. We herein report a case of MALS in an insulinoma patient.

Case Report

A 42-year-old man complaining of hypoglycemia was admitted to our hospital. On examination, his height, body weight, and body mass index were 165 cm, 62.8 kg, and 23.1, respectively. There was no abdominal pain or vomiting. His abdomen was soft with no tenderness, palpable mass, or organomegaly.

Laboratory data were as follows: serum glucose level 41 mg/dl, HbA1c 4.0%, and plasma insulin 7.7 μ U/mL. He had no significant medical history. He had never received insulin injection or medicine such as sulfonylurea compounds. Enhanced computed tomography revealed a 20-mm mass at the tail of pancreas (Fig. 1). He underwent a 72-h fast protocol that was terminated because of hypoglycemia. Finally, he was diagnosed as having insulinoma.

To determine the operability and location of the insulinoma, angiography and selective arterial calcium injection (SACI) test were performed. SACI test gave positive results for superior mesenteric artery (SMA). Angiography showed that blood flow in splenic artery was decreased. In addition, blood flow in SMA was increased. On the other hand, increase in blood flow in SMA was found from the dorsal pancreatic artery to insulinoma (Fig. 2A). In inferior

6-11-1 Omorinishi, Ota, Tokyo 143-8541, Japan

*Corresponding Author: tel: +81-3-3762-4151

e-mail: hiroshi.yoshino@med.toho-u.ac.jp

DOI: 10.14994/tohojmed.2018-012

Received May 21, 2018; Accepted July 26, 2018

Toho Journal of Medicine 5 (1), Mar. 1, 2019.

ISSN 2189-1990, CODEN: TJMOA2

pancreaticoduodenal artery (IPDA), retrograde blood flow was found from SMA to liver (Fig. 2B). Enhanced computed tomography revealed stenosis in the celiac artery (Fig. 3). Thus, he was diagnosed as suffering from MALS. Second SACI test gave positive results for dorsal pancreatic artery, but negative results for IPDA. Hence, the location of the insulinoma was determined. Laparoscopic spleen preserving distal pancreatectomy was successfully performed on the 97th day. Splenic artery and vein were divided and preserved (Fig. 4). Pathologic evaluation of the resected tissue revealed a well-differentiated tumor with a distinct border surrounding the exocrine pancreatic tissue. Immunohistochemical evaluation showed that the cells

were positive for CD56, synaptophysin, chromogranin-A, and insulin.

On post-operative day 10, the patient was found to have recovered without any surgical complication and was discharged.

Discussion

Insulinoma is the most common cause of hyperinsulinemic hypoglycemia. They are insulin-secreting tumors of pancreatic origin that cause hypoglycemia. Insulinoma had been found in 2-4 subjects per million, per year.²⁾ There are clinically significant variations of insulinoma, i.e., 4.8% of them demonstrate celiac artery stenosis.³⁾ According to




Fig. 1 Enhanced computed tomography revealed 20 mm of mass at the tail of pancreas.




Fig. 3 Enhanced computed tomography revealed stenosis in celiac artery (arrow).




Fig. 2

A) Angiogram reveals the blood flow from dorsal pancreatic artery to insulinoma (arrow).
 B) In inferior pancreaticoduodenal artery (IPDA), retrograde blood flow was found from SMA to liver by angiogram (black arrow).
 IPDA, inferior pancreaticoduodenal artery (white arrow); GDA, gastroduodenal artery.




Fig. 4 Tumor imaging during surgery

A) Mass at the tail of pancreas was clearly identified (arrow).

B) Resection of the tumor was successfully performed. Splenic artery and vein were divided and preserved (arrow).

Scott et al., diagnostic arteriography in combination with biochemical, pancreatic arterial anatomy, and regional perfusion data improved sensitivity of localization to 92.9%.³⁾ MALS typically occurs in young to middle-aged adults.⁴⁾ MALS has been reported in 4% of patients undergoing pancreaticoduodenectomy (PD).⁵⁾ In the present case, PD was not performed before Laparoscopic spleen preserving distal pancreatectomy. The incidence of celiac axis stenosis caused mainly by MALS is around 7.3% in asymptomatic individuals.⁶⁾ Decompression surgery, open surgery, laparoscopic surgery, and robotic surgery are hallmarks of the treatment.^{7, 8)}

To our knowledge, there were no reports of MALS in insulinoma patient. Because of MALS, celiac artery was compressed. Blood flow in splenic artery was decreased. In the present case, dorsal pancreatic artery was branched from SMA. On the other hand, blood flow in the dorsal pancreatic artery and SMA were increased. According to Woodburne, 21.5% of dorsal pancreatic artery arises close to the source of the SMA.⁹⁾ Therefore, blood flow from dorsal pancreatic artery to insulinoma was found. In addition, retrograde blood flow was found from SMA to liver in IPDA. MALS should be considered when angiography of the area surrounding the pancreas reveals blood flow different from that observed in normal anatomy.

Acknowledgements: We thank Dr. Masaru Tsuchiya, (Department of Surgery, Division of Gastroenterological Surgery) for helpful discussion.

Consent: Informed consent was obtained from the patient and

family for publication of this case report and the accompanying images.

Conflicts of interest: None declared.

References

- 1) Harjola PT. A rare obstruction of the coeliac artery. *Ann Chir Gynaecol Fenn.* 1963; 52: 547-50.
- 2) Oberg K, Eriksson B. Endocrine tumors of the pancreas. *Best Pract Res Clin Gastroenterol.* 2005; 19: 753-81.
- 3) Thompson SM, Vella A, Service F, Grant CS, Thompson GB, Andrews JC. Impact of variant pancreatic arterial anatomy and overlap in regional perfusion on the interpretation of selective arterial calcium stimulation with hepatic venous sampling for preoperative localization of occult insulinoma. *Surgery.* 2015; 158: 162-72.
- 4) Schweizer P, Berger S, Schweizer M, Schaefer J, Beck O. Arcuate ligament vascular compression syndrome in infants and children. *J Pediatr Surg.* 2005; 40: 1616-22.
- 5) Farma JM, Hoffman JP. Neoplastic celiac axis occlusion in patients undergoing pancreaticoduodenectomy. *Am J Surg.* 2007; 193: 341-4.
- 6) Park CM, Chung JW, Kim HB, Shin SJ, Park JH. Celiac axis stenosis: incidence and etiologies in asymptomatic individuals. *Korean J Radiol.* 2001; 21: 8-13.
- 7) Tulloch AW, Jimenez JC, Lawrence PF, Dutson EP, Moore WS, Rigberg DA, et al. Laparoscopic versus open celiac ganglionectomy in patients with median arcuate ligament syndrome. *J Vasc Surg.* 2010; 52: 1283-9.
- 8) Do MV, Smith TA, Bazan HA, Sternbergh WC 3rd, Abbas AE, Richardson WS. Laparoscopic versus robot-assisted surgery for median arcuate ligament syndrome. *Surg Endosc.* 2013; 27: 4060-6.
- 9) Woodburne RT, Olsen LL. The arteries of the pancreas. *Anat Rec.* 1951; 111: 255-70.

Toho Journal of Medicine. Toho Journal of Medicine is an Open Access journal distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view the details of this license, please visit (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).