

東邦大学学術リポジトリ

Toho University Academic Repository

タイトル	Association of Repeated Defibrillation with Outcomes for Out of Hospital Cardiac Arrest Associated with Ventricular Fibrillation
作成者（著者）	Takaya, Tsubota / Mitsuru, Honda / Katsunori, Yoshihara / Takanori, Ikeda
公開者	The Medical Society of Toho University
発行日	2016.06
ISSN	21891990
掲載情報	Toho Journal of Medicine. 2(2). p.53 60.
資料種別	学術雑誌論文
内容記述	Original Article
著者版フラグ	publisher
JaLCDOI	info:doi/10.14994/tohojmed.2015.016
メタデータのURL	https://mylibrary.toho u.ac.jp/webopac/TD59677828

Association of Repeated Defibrillation with Outcomes for Out-of-Hospital Cardiac Arrest Associated with Ventricular Fibrillation

Takaya Tsubota^{1)*} Mitsuru Honda²⁾ Katsunori Yoshihara³⁾
and Takanori Ikeda¹⁾

¹⁾Division of Cardiovascular Medicine (Omori), Department of Internal Medicine,
School of Medicine, Faculty of Medicine, Toho University

²⁾Department of General Medicine and Emergency Care, School of Medicine, Faculty of Medicine, Toho University

³⁾Department of Emergency and Disaster Medicine, Toho University Omori Medical Center

ABSTRACT

Background: Ventricular fibrillation (VF) is a life-threatening arrhythmia, and the success rate for defibrillation decreases rapidly with increasing time since collapse. Chest compression must be interrupted during rhythm analysis and defibrillation, and this may adversely affect outcomes. This study investigated the effectiveness of repeated defibrillation for persons with cardiogenic out-of-hospital cardiac arrest (OHCA) associated with VF.

Methods: Using Utstein template data collected throughout Japan during the period from July 2006 through December 2010, we identified patients with VF on an initial electrocardiogram who underwent defibrillation within 10 minutes of collapse (N=9865). To determine the optimal frequency of defibrillation, we evaluated rates of return of spontaneous circulation (ROSC) and survival at 1 month with a good neurologic outcome (cerebral performance category [CPC] 1–2).

Results: ROSC was achieved in 38.2% of patients, and 28.5% had a CPC score of 1 or 2 at 1 month post-event. Receiver operating characteristic curve analysis showed that a defibrillation frequency cutoff value of 1.5 ($p < 0.001$; 95% confidence interval, 0.61–0.63) was optimal for ROSC. The same cutoff was associated with a CPC of 1 to 2 ($p < 0.001$; 95% confidence interval, 0.57–0.60). As the number of shocks increased, the rates of ROSC and good neurologic outcomes decreased. Among patients who achieved ROSC and a good neurologic outcome, half had received 1 defibrillation.

Conclusions: Repeated defibrillation may be associated with worse outcomes.

Toho J Med 2 (2): 53–60, 2016

KEYWORDS: defibrillation, ventricular fibrillation (VF), cardiac arrest, out-of-hospital

Improving the rate of survival after out-of-hospital cardiac arrest (OHCA) is an important issue in emergency

medicine and intensive care. A number of randomized studies of rates of successful resuscitation have high-

1, 2, 3) 6–11–1 Omorinishi, Ota, Tokyo 143–8541, Japan

*Corresponding Author: tel: +81-(0)3-3762-4151

e-mail: tsubota@med.toho-u.ac.jp

DOI: 10.14994/tohojmed.2015.016

Received Sept. 29, 2015; Accepted Apr. 23, 2016

Toho Journal of Medicine 2 (2), June 1, 2016.

ISSN 2189–1990, CODEN: TJMOA2

highlighted the importance of early access to emergency medical services (EMS), cardiopulmonary resuscitation (CPR), and defibrillation, as well as the effectiveness of advanced cardiac life support and integrated treatment after return of spontaneous circulation (ROSC), in improving survival after OHCA. One-month survival and favorable neurologic outcomes after witnessed OHCA increased from 5% and 2%, respectively, in 1998 to 12% and 6% in 2006, but these rates are nevertheless low.^{1,2)}

Most OHCA involve adults, and cardiac arrest associated with ventricular fibrillation (VF) and pulseless ventricular tachycardia (VT) has the best resuscitation rates.³⁾ VF accounts for approximately 20% of all OHCA, and the elapsed time between cardiac arrest and electrocardiogram (ECG) recording is less than 3 minutes in 50% to 60% of cases.⁴⁾ The outcomes for patients with VF or pulseless VT cardiac arrest are far better than those for patients with asystole or pulseless electrical activity cardiac arrest,^{3,5,6)} and the rates of 1-month survival and favorable neurologic outcomes in witnessed cardiogenic OHCA for which the initial rhythm was VF increased from 9.8% in 2005 to 20.6% in 2009.⁷⁾

Continuous chest compressions and early defibrillation are important considerations in VF. Shorter pre-shock pauses are also associated with successful defibrillation (odds ratio [OR], 1.86 per 5-second decrease).⁸⁾ Patients with VF who receive immediate CPR from bystanders can survive without neurologic sequelae, provided that defibrillation is performed within 5 to 10 minutes.^{9,10)} However, interrupting chest compressions during ECG interpretation or defibrillation may worsen outcomes.^{11,12)} VF is a life-threatening arrhythmia, and the rate of successful defibrillation decreases over time¹²⁾; therefore, early defibrillation is important. In 2010, the mean elapsed time between contact with EMS and the arrival of emergency personnel to assist the patient was 8.1 minutes in Japan.¹³⁾ However, studies have not addressed the association between frequency of defibrillation and outcomes. This study investigated the association of number of EMS-delivered on-site defibrillations with 1-month survival and neurologic outcomes.

Methods

Study design and setting

A nationwide population-based registry of OHCA throughout Japan is maintained by the Fire and Disaster Management Agency. Using this registry, we prospec-

tively collected Utstein template data for the period from July 1, 2006 through December 31, 2010^{14,15)} The subjects of this observational study were persons with a witnessed OHCA who received CPR from EMS personnel. The sample was further limited to those patients with VF during their initial ECG who received a first defibrillation by EMS personnel within 10 minutes of their witnessed cardiogenic OHCA. This study was performed in accordance with Japanese CPR guidelines, which are based on the 2005 American Heart Association (AHA) guidelines, as a 1-shock protocol, that is to say when VF or pulseless VT rhythm was present, 1-shock was delivered and CPR was immediately resumed, beginning with chest compressions.¹⁶⁾

Cardiac arrest was classified as cardiogenic if it was not endogenous (caused by a central nervous system disorder, circulatory disorder, malignant tumor, or other similar condition), exogenous (caused by, for example, trauma, strangulation, drowning, or drug overdose), or attributable to disorders believed to be noncardiogenic. Cardiogenic and noncardiogenic cardiac arrests were distinguished on the basis of the clinical judgment of the doctor who provided initial care after the patient arrived at the hospital.

The Resuscitation Science Group Subcommittee of the Japanese Circulation Society was provided with data from the governmental legal procedures. This study was reviewed and approved by the ethics committee of Toho University (approval number 25086).

EMS system

Data published by the Japanese Fire and Disaster Management Agency in 2010 showed that EMS were contacted approximately 5.46 million times; the mean interval between receiving a call and arriving at the patient's side was 8.1 minutes.¹³⁾ In Japan, an EMS team typically includes 3 emergency providers, including at least 1 emergency lifesaving technician. Emergency lifesaving technicians are permitted to insert intravenous lines and adjunct airways and to use semiautomatic defibrillators before OHCA patients arrive at a hospital.

Since July 2004, trained emergency medical technicians have been allowed to perform tracheal intubation and, in 2006, they were authorized to administer intravenous adrenaline. All emergency technicians can perform CPR according to the Japanese guidelines, which are based on the 2005 AHA guidelines. Emergency lifesaving technicians are not permitted to terminate CPR on-site, and most patients are transported to a hospital.¹⁶⁾

Data collection

The OHCA data included information on age, sex, presence of a witness, whether and how CPR was performed by a bystander, ECG waveform classification, how the airway was secured, drugs administered, whether and how many times defibrillation was performed, whether ROSC was achieved, and when these actions were performed. In addition, the times at which EMS personnel were contacted, arrived at the patient's side, and arrived at the hospital with the patient were also recorded. To assess outcomes, 1-month survival and neurologic outcome, as determined by cerebral performance category (CPC),¹⁷⁾ were recorded. Patient CPC was determined by the attending physician. A good neurologic outcome was defined as a CPC score of 1 or 2 and a poor outcome as a CPC score of 3 to 5. Successful defibrillation was defined as achieving ROSC.

Statistical analysis

We compared patients who achieved ROSC with those who did not (ROSC and non-ROSC groups), and those with good and poor neurologic outcomes. Multivariate analysis using logistic regression models (forward selection) was used to examine factors associated with ROSC and a CPC score of 1 or 2; adjusted ORs and 95% confidence intervals (CIs) were calculated. The optimal number of defibrillations needed to achieve ROSC and a CPC score of 1 or 2 were determined using receiver operating characteristic (ROC) curves to calculate cutoff points, which were defined as the maximum point calculated using the equation sensitivity + specificity. The SPSS version 20 software package (International Business Machines [IBM] Corp., Armonk, NY, USA) was used to perform the analyses, and all values are expressed as mean \pm SD or number and percentage. Intergroup comparisons were done by using the *t* test or χ^2 test. A *p* value of less than 0.05 was considered to indicate statistical significance.

Results

In total, 510727 OHCA (281453 cardiogenic and 229274 noncardiogenic) were recorded between July 1, 2006 and December 31, 2010. Among the cardiogenic OHCA, 114815 were witnessed; 9865 of these occurred in persons with an initial ECG waveform of VF who received initial defibrillation by EMS within 10 minutes of cardiogenic cardiac arrest (Fig. 1).

Baseline characteristics

Among the 9865 patients included in the study, 3812 had




Fig. 1 Study flow for out of hospital cardiac arrest patients.

VF: ventricular fibrillation, EMS: emergency medical service

circulation restored and 6053 did not (Table 1). Among the 3812 patients with restored circulation, 2811 had a good neurologic outcome (CPC score, 1–2) and 7054 had a poor outcome (CPC score, 3–5) at 1 month. Younger patients were significantly more likely to have circulation restored and to have a good neurologic outcome. Similarly, early CPR, early defibrillation, chest compression and ventilation, and use of a biphasic waveform defibrillator were significantly associated with restoration of circulation and a good neurologic outcome. Use of an advanced airway device and use of adrenaline by EMS personnel were significantly more frequent in non-ROSC patients and those with a CPC score of 3 to 5. The time taken for EMS personnel to arrive at the patient's side was not associated with ROSC or a good neurologic outcome. The interval between collapse and hospital arrival was longer in the ROSC group and shorter in patients with a good neurologic outcome.

Multivariate analysis of ROSC

Multivariate analysis (Table 2) showed that ROSC was associated with chest compression (adjusted OR, 1.293; 95% CI, 1.167-1.433) and use of a biphasic waveform defibrillator by EMS personnel (adjusted OR, 1.208; 95% CI, 1.078-1.354). ROSC was associated with early defibrillation (adjusted OR per 1-minute increment between collapse and defibrillation time, 0.942; 95% CI, 0.928-0.956) and age (adjusted OR per 1-year increase in age, 0.989; 95% CI, 0.986-0.992). Use of an advanced airway device was in-

Table 1 Baseline characteristics of patients

	ROSC N = 3812 (38.2%)	Non-ROSC N = 6053 (61.8%)	P value	CPC 1-2 N = 2811 (28.5%)	CPC 3-5 N = 7054 (71.5%)	P value
Age, years	63.4 ± 15.2	65.2 ± 15.4	<0.001	60.6 ± 15.0	66.0 ± 15.2	<0.001
Men (%)	2985 (78.3)	4861 (80.3)	0.014	2272 (80.8)	55742 (79.0)	0.054
Bystander CPR						
Chest compression (%)	1534 (40.2)	2306 (38.0)	0.029	1182 (41.9)	2658 (37.7)	0.001
Ventilation (%)	530 (13.8)	746 (12.3)	0.039	410 (14.5)	866 (12.3)	0.003
Biphasic waveform defibrillation (%)	3207 (84.1)	4972 (82.1)	0.013	2422 (86.1)	5757 (81.6)	<0.001
Number of defibrillations	1.9 ± 1.3	2.5 ± 1.7	<0.001	1.9 ± 1.3	2.4 ± 1.6	<0.001
Use of advanced airway device (%)	960 (25.1)	2800 (46.3)	<0.001	568 (20.1)	3192 (45.3)	<0.001
Use of adrenaline by EMS (%)	318 (8.2)	853 (14.1)	<0.001	145 (5.1)	1026 (14.5)	<0.001
Interval from collapse to EMS arrival at patient's side (min)	5.5 ± 2.5	5.6 ± 2.5	0.101	5.5 ± 2.5	5.6 ± 2.6	0.08
Interval from collapse to start of CPR (min)	4.5 ± 3.2	5.0 ± 3.1	<0.001	4.4 ± 3.2	5.0 ± 3.1	<0.001
Interval from collapse to defibrillation (min)	5.8 ± 3.3	6.6 ± 3.0	<0.001	5.7 ± 3.3	6.5 ± 3.1	<0.001
Interval from collapse to hospital arrival (min)	26.6 ± 14.1	25.9 ± 11.3	0.006	25.0 ± 14.4	26.6 ± 11.6	<0.001

ROSC: return of spontaneous circulation, CPC: cerebral performance category, CPR: cardiopulmonary resuscitation, EMS: emergency medical services

Table 2 Logistic regression analysis of the associations of patient and treatment characteristics with return of spontaneous circulation

Variable	Adjusted odds ratio (95% CI)	P value
Age * ¹	0.989 (0.986-0.992)	<0.001
Male, sex	0.860 (0.773-0.958)	0.006
Chest compression	1.293 (1.167-1.433)	<0.001
Ventilation (%)	1.050 (0.913-1.209)	0.492
Biphasic waveform defibrillation	1.208 (1.078-1.354)	0.001
Number of defibrillations * ²	0.753 (0.729-0.779)	0.001
Use of advanced airway device	0.446 (0.406-0.490)	<0.001
Use of adrenaline by EMS	0.938 (0.810-1.087)	0.396
Interval from collapse to defibrillation * ³	0.942 (0.928-0.956)	<0.001

CI: confidence interval, CPR: cardiopulmonary resuscitation, EMS: emergency medical service
*¹: per 1-year increment, *²: per additional defibrillation, *³: per 1-min increment

versely associated with ROSC (adjusted OR, 0.446; 95% CI, 0.406-0.490).

Multivariate analysis of good neurologic outcome

Multivariate analysis (Table 3) showed that a good neurologic outcome was associated with chest compression (adjusted OR, 1.494; 95% CI, 1.334-1.673) and use of a biphasic waveform defibrillator by EMS personnel (1.492; 1.311-1.698). ROSC was associated with early defibrillation (adjusted OR per 1-minute increment, 0.922; 95% CI, 0.907-0.936) and age (adjusted OR per 1-year increment, 0.974;

95% CI, 0.971-0.977). As was the case for factors contributing to ROSC, however, a good neurologic outcome was not associated with use of an advanced airway device (adjusted OR, 0.361; 95% CI, 0.324-0.402) or adrenaline (0.497; 0.410-0.602).

Number of shocks

As the number of shocks increased, the rates of ROSC and good neurologic outcomes decreased (Table 4). Among patients who achieved ROSC and a good neurologic outcome, half had received 1 defibrillation.

Table 3 Logistic regression analysis of the associations of patient and treatment characteristics with a good neurologic outcome

Variable	Adjusted odds ratio (95% CI)	P value
Age * ¹	0.974 (0.971-0.977)	<0.001
Male sex	0.986 (0.875-1.111)	0.816
Chest compression	1.494 (1.334-1.673)	<0.001
Ventilation (%)	1.041 (0.895-1.211)	0.603
Biphasic waveform defibrillation	1.492 (1.311-1.698)	<0.001
Number of defibrillations * ²	0.829 (0.801-0.859)	<0.001
Use of advanced airway device	0.361 (0.324-0.402)	<0.001
Use of adrenaline by EMS	0.497 (0.410-0.602)	<0.001
Interval from collapse to defibrillation * ³	0.922 (0.907-0.936)	<0.001

CI: confidence interval, CPR: cardiopulmonary resuscitation, EMS: emergency medical service

*¹: per 1-year increment; *²: per additional defibrillation; *³: per 1-min increment

Table 4 ROSC and CPC in relation to the number of shocks administered

Number of shocks	ROSC (N = 3812)	Non-ROSC (N = 6053)	CPC 1-2 (N = 2811)	CPC 3-4 (N = 7054)
1 (%)	2044 (53.6)	2104 (32.4)	1455 (51.8)	2693 (38.2)
2 (%)	942 (24.7)	1567 (24.1)	691 (24.6)	1818 (25.8)
3 (%)	460 (12.1)	1085 (16.7)	369 (13.1)	1176 (16.7)
4 (%)	192 (5.0)	565 (8.7)	140 (5.0)	617 (8.7)
≥5 (%)	174 (4.6)	732 (12.1)	156 (5.5)	750 (10.6)

ROSC: return of spontaneous circulation, CPC: cerebral performance category

ROC curve analysis

The ROC curve analysis of ROSC and number of defibrillations showed that the cutoff value for the optimal number of shocks required to maximize successful ROSC was 1.5, with a sensitivity of 65%, a specificity of 54%, and an area under the curve (AUC) of 0.62 (p<0.001; 95% CI, 0.61-0.63) (Fig. 2A). A similar analysis showed that the optimal number of defibrillations required to maximize neurologic outcome (CPC score, 1-2) was also 1.5 (sensitivity, 62%; specificity, 52%; AUC, 0.59; p<0.05; 95% CI, 0.57-0.60) (Fig. 2B).

Time to hospital arrival

The mean time between collapse and hospital arrival was 24.9±12.6 minutes for patients receiving 1 or 2 shocks and 28.7±11.9 minutes for those receiving 3 or more shocks. The time between collapse and hospital arrival was significantly shorter for patients receiving 1 or 2 shocks than for those receiving 3 or more shocks (p<0.001).

Discussion

ROC curve analysis showed that the optimal number of pre-hospital shocks for OHCA patients (as determined by rates of ROSC and a CPC score of 1-2) was 1.5; thus, use of 2 shocks is appropriate in clinical practice. As the number of shocks increased, rates of ROSC and good neurologic outcomes decreased, and about half of patients were successfully treated with 1 shock. If frequent shocks are administered, EMS personnel must stop the ambulance and interrupt chest compressions for each cycle of ECG analysis and shock administration. Coronary perfusion pressure decreases if chest compressions are interrupted, causing a decline in coronary arteries. Even when chest compression is restarted, several consecutive compressions are required for coronary artery perfusion to return to its previous level.¹⁸⁾ Interrupting chest compressions during CPR was found to reduce ROSC rate, survival rate, and post-resuscitation cardiac function.¹⁹⁾ Minimizing the chest compression interruptions necessary to analyze cardiac rhythm, defibrillate, and administer advanced life support




Fig. 2 Receiver operating curve (ROC) analysis of the number of defibrillations and return of spontaneous circulation (A) and a good neurologic outcome (B). A cerebral performance category (CPC) score of 1 or 2 was defined as a good neurologic outcome.

AUC: area under the curve, CI: confidence interval

improves defibrillation outcomes.^{20, 21)}

VF is divided into 3 stages according to the interval from onset. The period within 4 to 5 minutes of VF onset is known as the electrical phase and is characterized by mild ischemia. ROSC can easily be obtained by rapid electrical defibrillation during this phase. The period 4 to 10 minutes after VF onset is referred to as the hemodynamic or circulatory phase. During this phase, myocardial adenosine triphosphate is depleted, which increases the difficulty of restarting a spontaneous rhythm, even if defibrillation is performed. Performing chest compressions during this phase maintains cerebral and coronary artery perfusion and reduces ischemia, which increases the possibility of achieving ROSC by defibrillation. The metabolic phase begins at 10 minutes after VF onset. During this phase, ROSC is difficult to achieve by regular CPR, including defibrillation; treatment of the underlying cause or adjunctive treatment is required.²²⁾

The outcomes for shock-resistant VF are disappointing: ROSC in 20.4% of cases and a CPC score of 1 or 2 in 5.6% of cases in 2006.²³⁾ More time passes as additional unsuccessful defibrillations are performed, and the success rate decreases. The need for multiple defibrillation attempts increases interruptions to chest compression; thus, unsuccessful defibrillations result in myocardial damage and decreased survival.^{24, 25)}

The time taken for EMS personnel to arrive at the patient's side was not associated with ROSC or a good neu-

rologic outcome; however, CPR and defibrillation should be performed as quickly as possible, and the patient should be hospitalized quickly. For persons with shock-resistant VF, early transportation for administration of anti-arrhythmic drugs, emergency cardiopulmonary bypass, and therapeutic hypothermia may be more important than repeated defibrillation.²⁶⁾ To improve neurologic outcomes, the frequency of defibrillation — namely, limiting the number of shocks — should be considered by EMS personnel on the scene.

Bystander CPR, especially chest compression, was an important factor in achieving ROSC and good neurologic outcomes. Previous studies reported that a longer interval between cardiac arrest and defibrillation was associated with worse patient survival.²⁷⁾ If CPR is withheld, the survival rate after witnessed VF cardiac arrest declines by an average of 7% to 10% for each minute between collapse and defibrillation.¹²⁾ If CPR is performed by bystanders, however, the survival rate declines by only 3% to 4% per minute.^{12, 27-29)} Performing CPR also reportedly doubles or triples the survival rate for a patient with a witnessed cardiac arrest, even if some time elapses before defibrillation.^{12, 30, 31)}

Our results suggest that adrenaline administration and use of advanced airway devices is disadvantageous to ROSC and worsens neurologic outcomes. However, adrenaline and use of advanced airway devices are not indicated for patients in whom ROSC was achieved by de-

fibrillation and thus may only have been used for patients without ROSC in the present study. In other words, only patients in worse condition received an advanced airway device.

Use of a biphasic defibrillator was an important factor in achieving ROSC and a good neurologic outcome in the present patients. Biphasic defibrillators are reported to have a higher defibrillation rate than monophasic defibrillators.³²⁻³⁴ The considerable energy needed for a successful shock causes myocardial damage. Biphasic defibrillators use lower energy than monophasic defibrillators and are less likely to cause post-shock VF.³⁵

As compared with men, women had a higher rate of ROSC, but the proportions of men and women with a good neurologic outcome did not significantly differ. Some studies reported that the central nervous system and heart of women are protected by the effect of endogenous female hormones.³⁶ An animal study found that hormone administration improved cardiac arrest outcomes.³⁷ Studies clearly show that rates of survival and good neurologic outcomes decrease with advancing age.³⁸ In this study, the men were younger (mean age: 62.7 ± 14.5 years for men and 71.2 ± 16.6 years for women, $p < 0.005$). Thus, age appears to be strongly associated with neurologic outcome.

Study limitations

This study had some limitations. First, it was not a randomized controlled study. Second, we only analyzed pre-hospitalization data and thus have no information on treatments received in hospital, anti-arrhythmic drug use, coronary interventions, emergency cardiopulmonary bypass, or use of therapeutic hypothermia. Neurologic outcomes would likely be improved by a combination of emergency cardiopulmonary bypass, percutaneous coronary intervention, and early therapeutic hypothermia for OHCA patients unresponsive to conventional CPR for whom the period of circulatory arrest is comparatively short and the cause of cardiac arrest is known.²⁶ However, the evidence for this hypothesis is not conclusive.

Conclusions

Bystander CPR, biphasic waveform defibrillation, and temporal factors were important in achieving ROSC and good neurologic outcomes. Analysis of the data showed that, for an OHCA patient with VF, 2 shocks was the optimal number of defibrillations by EMS personnel on the scene. These data suggest that repeated defibrillation is associated with worse prognosis and that patients should

be transported to a hospital as quickly as possible while chest compressions are maintained.

We would like to thank the EMS personnel, the Fire and Disaster Management Agency, and the physicians involved in establishing and maintaining the Utstein database.

Conflicts of interest: The authors have no conflicts of interest to disclose.

References

- 1) Field JM, Hazinski MF, Sayre MR, Chameides L, Schexnayder SM, Hemphill R, et al. Part 1: executive summary: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*. 2010; 122 Suppl 3: S640-56.
- 2) Iwami T, Nichol G, Hiraide A, Hayashi Y, Nishiuchi T, Kajino K, et al. Continuous improvements in "chain of survival" increased survival after out-of-hospital cardiac arrests: a large-scale population-based study. *Circulation*. 2009; 119: 728-34.
- 3) Rea TD, Cook AJ, Stiell IG, Powell J, Bigham B, Callaway CW, et al; Resuscitation Outcomes Consortium Investigators. Predicting survival after out-of-hospital cardiac arrest: role of the Utstein data elements. *Ann Emerg Med*. 2010; 55: 249-57.
- 4) SOS-KANTO Committee. Incidence of ventricular fibrillation in patients with out-of-hospital cardiac arrest in Japan: survey of survivors after out-of-hospital cardiac arrest in Kanto area (SOS-KANTO). *Circ J*. 2005; 69: 1157-62.
- 5) Nadkarni VM, Larkin GL, Peberdy MA, Carey SM, Kaye W, Mancini ME, et al; National Registry of Cardiopulmonary Resuscitation Investigators. First documented rhythm and clinical outcome from in-hospital cardiac arrest among children and adults. *JAMA*. 2006; 295: 50-7.
- 6) Meaney PA, Nadkarni VM, Kern KB, Indik JH, Halperin HR, Berg RA. Rhythms and outcomes of adult in-hospital cardiac arrest. *Crit Care Med*. 2010; 38: 101-8.
- 7) Kitamura T, Iwami T, Kawamura T, Nitta M, Nagao K, Nonogi H, et al. Japanese Circulation Society Resuscitation Science Study Group. Nationwide improvements in survival from out-of-hospital cardiac arrest in Japan. *Circulation*. 2012; 126: 2834-43.
- 8) Edelson DP, Abella BS, Kramer-Johansen J, Wik L, Myklebust H, Barry AM, et al. Effects of compression depth and pre-shock pauses predict defibrillation failure during cardiac arrest. *Resuscitation*. 2006; 71: 137-45.
- 9) Wik L, Hansen TB, Fylling F, Steen T, Vaagenes P, Auestad BH, et al. Delaying defibrillation to give basic cardiopulmonary resuscitation to patients with out-of-hospital ventricular fibrillation: a randomized trial. *JAMA*. 2003; 289: 1389-95.
- 10) Cobb LA, Fahrenbruch CE, Walsh TR, Copass MK, Olsufka M, Breskin M, et al. Influence of cardiopulmonary resuscitation prior to defibrillation in patients with out-of-hospital ventricular fibrillation. *JAMA*. 1999; 281: 1182-8.
- 11) Christenson J, Andrusiek D, Everson-Stewart S, Kudenchuk P, Hostler D, Powell J, et al. Chest compression fraction determines survival in patients with out-of-hospital ventricular fibrillation. *Circulation*. 2009; 120: 1241-7.
- 12) Larsen MP, Eisenberg MS, Cummins RO, Hallstrom AP. Predict-

- ing survival from out-of-hospital cardiac arrest: a graphic model. *Ann Emerg Med.* 1993; 22: 1652-8.
- 13) Ambulance Service Planning Office [Internet]. Tokyo: Fire and Disaster Management Agency (Japan) [Cited 2013 June 24] Available from: http://www.fdma.go.jp/neuter/topics/kyukyuk_yujo_genkyo/h24/01_kyukyuu.pdf
 - 14) Cummins RO, Chamberlain DA, Abramson NS, Allen M, Baskett PJ, Becker L, et al. Recommended guideline for uniform reporting of data from out-of-hospital cardiac arrest: the Utstein Style: A statement for health professionals from a task force of the American Heart Association, the European Resuscitation Council, the Heart and Stroke Foundation of Canada, and the Australian Resuscitation Council. *Circulation.* 1991; 84: 960-75.
 - 15) Jacobs I, Nadkarni V, Bahr J, Berg RA, Billi JE, Bossaert L, et al; ILCOR Task Force on Cardiac Arrest and Cardiopulmonary Resuscitation Outcomes. Cardiac arrest and cardiopulmonary resuscitation outcome reports: update and simplification of the Utstein templates for resuscitation registries. A statement for healthcare professionals from a task force of the International Liaison Committee on Resuscitation (American Heart Association, European Resuscitation Council, Australian Resuscitation Council, New Zealand Resuscitation Council, Heart and Stroke Foundation of Canada, InterAmerican Heart Foundation, Resuscitation Council of Southern Africa). *Circulation.* 2004; 110: 3385-97.
 - 16) Resuscitation and Emergency Care Guidelines Committee of Japan Foundation for Emergency Medicine. *Kyukyu sosei ho no shishin 2005 iryojujisha yo.* 3rd ed. Tokyo: Herusu Shuppan; 2007. Japanese.
 - 17) Jennett B, Bond M. Assessment of outcome after severe brain damage. *Lancet.* 1975; 1: 480-4.
 - 18) Berg RA, Sanders AB, Kern KB, Hilwig RW, Heidenreich JW, Porter ME, et al. Adverse hemodynamic effects of interrupting chest compressions for rescue breathing during cardiopulmonary resuscitation for ventricular fibrillation cardiac arrest. *Circulation.* 2001; 104: 2465-70.
 - 19) Eftestøl T, Sunde K, Steen PA. Effects of interrupting precordial compressions on the calculated probability of defibrillation success during out-of-hospital cardiac arrest. *Circulation.* 2002; 105: 2270-3.
 - 20) Rea TD, Helbock M, Perry S, Garcia M, Cloyd D, Becker L, et al. Increasing use of cardiopulmonary resuscitation during out-of-hospital ventricular fibrillation arrest: survival implications of guideline changes. *Circulation.* 2006; 114: 2760-5.
 - 21) Hinchey PR, Myers JB, Lewis R, De Maio VJ, Reyer E, Licatase D, et al. Improved out-of-hospital cardiac arrest survival after the sequential implementation of 2005 AHA guidelines for compressions, ventilations, and induced hypothermia: the Wake County experience. *Ann Emerg Med.* 2010; 56: 348-57.
 - 22) Xie J, Weil MH, Sun S, Tang W, Sato Y, Jin X, et al. High-energy defibrillation increases the severity of postresuscitation myocardial dysfunction. *Circulation.* 1997; 96: 683-8.
 - 23) Yamaguchi H, Wei MH, Tang W, Kamohara T, Jin X, Bisera J. Myocardial dysfunction after electrical defibrillation. *Resuscitation.* 2002; 54: 289-96.
 - 24) Weisfeldt ML, Becker LB. Resuscitation after cardiac arrest: a 3-phase time-sensitive model. *JAMA.* 2002; 288: 3035-8.
 - 25) Sakai T, Iwami T, Tasaki O, Kawamura T, Hayashi Y, Rinka H, et al. Incidence and outcomes out-of-hospital cardiac arrest with shock-resistant ventricular fibrillation: Data from a large population-based cohort. *Resuscitation.* 2010; 81: 956-61.
 - 26) Nagao K, Kikushima K, Watanabe K, Tachibana E, Tominaga Y, Tada K, et al. Early induction of hypothermia during cardiac arrest improves neurological outcomes in patients with out-of-hospital cardiac arrest who undergo emergency cardiopulmonary bypass and percutaneous coronary intervention. *Circ J.* 2010; 74: 77-85.
 - 27) Chan PS, Krumholz HM, Nichol G, Nallamothu BK; American Heart Association National Registry of Cardiopulmonary Resuscitation Investigators. Delayed time to defibrillation after in-hospital cardiac arrest. *N Engl J Med.* 2008; 358: 9-17.
 - 28) Valenzuela TD, Roe DJ, Cretin S, Spaite DW, Larsen MP. Estimating effectiveness of cardiac arrest interventions: a logistic regression survival model. *Circulation.* 1997; 96: 3308-13.
 - 29) Stiell IG, Wells GA, Field B, Spaite DW, Nesbitt LP, De Maio VJ, et al; Ontario Prehospital Advances Life Support Study Group. Advanced cardiac life support in out-of-hospital cardiac arrest. *N Engl J Med.* 2004; 351: 647-56.
 - 30) Swor RA, Jackson RE, Cynar M, Sadler E, Basse E, Boji B, et al. Bystander CPR, ventricular fibrillation, and survival in witnessed, unmonitored out-of-hospital cardiac arrest. *Ann Emerg Med.* 1995; 25: 780-4.
 - 31) Holmberg M, Holmberg S, Herlitz J. Incidence, duration and survival of ventricular fibrillation in out-of-hospital cardiac arrest patients in Sweden. *Resuscitation.* 2000; 44: 7-17.
 - 32) Schneider T, Martens PR, Paschen H, Kuisma M, Wolcke B, Gliner BE, et al. Multicenter, randomized, controlled trial of 150-J biphasic shocks compared with 200- to 360-J monophasic shocks in the resuscitation of out-of-hospital cardiac arrest victims. Optimized Response to Cardiac Arrest (ORCA) Investigators. *Circulation.* 2000; 102: 1780-7.
 - 33) Morrison LJ, Dorian P, Long J, Vermeulen M, Schwartz B, Sawadsky B, et al. Out-of-hospital cardiac arrest rectilinear biphasic to monophasic damped sine defibrillation waveforms with advanced life support intervention trial (ORBIT). *Resuscitation.* 2005; 66: 149-57.
 - 34) van Alem AP, Chapman FW, Lank P, Hart AA, Koster RW. A prospective, randomized and blinded comparison of first shock success of monophasic and biphasic waveforms in out-of-hospital cardiac arrest. *Resuscitation.* 2003; 58: 17-24.
 - 35) Kudenchuk PJ, Cobb LA, Copass MK, Olsufka M, Maynard C, Nichol G. Transthoracic incremental monophasic versus biphasic defibrillation by emergency responders (TIMBER): a randomized comparison of monophasic with biphasic waveform ascending energy defibrillation for the resuscitation of out-of-hospital cardiac arrest due to ventricular fibrillation. *Circulation.* 2006; 114: 2010-8.
 - 36) Mendelsohn ME, Karas RH. Molecular and cellular basis of cardiovascular gender differences. *Science.* 2005; 308: 1583-7.
 - 37) Jover T, Tanaka H, Calderone A, Oguro K, Bennett MV, Etgen AM, et al. Estrogen protects against global ischemia-induced neuronal death and prevents activation of apoptotic signaling cascades in the hippocampal CA1. *J Neurosci.* 2002; 22: 2115-24.
 - 38) Iwami T, Hiraide T, Nakanishi N, Hayashi Y, Nishiuchi T, Yukioka H, et al. Age and sex analyses of out-of-hospital cardiac arrest in Osaka, Japan. *Resuscitation.* 2003; 57: 145-52.