

# 東邦大学学術リポジトリ

Toho University Academic Repository

タイトル	The growth factor midkine may play a pathophysiological role in rheumatoid arthritis
別タイトル	成長因子ミッドカインは関節リウマチの病態形成に關与する
作成者(著者)	進藤, 恵実子
公開者	東邦大学
発行日	2016.03
掲載情報	東邦大学大学院医学研究科 博士論文. 4.
資料種別	学位論文
内容記述	主査: 三上哲夫 / タイトル: The growth factor midkine may play a pathophysiological role in rheumatoid arthritis / 著者: Emiko Shindo, Toshihiro Nanki, Natsuko Kusunoki, Kotaro Shikano, Mai Kawazoe, Hiroshi Sato, Kaichi Kaneko, Sei Muraoka, Makoto Kaburaki, Yoshikiyo Akasaka, Hideaki Shimada, Tomoko Hasunuma, Shinichi Kawai / 掲載誌: Modern Rheumatology / 巻号・発行年等: 27(1):54-59, 2017 / 本文ファイル: 査読後原稿
著者版フラグ	ETD
報告番号	32661甲第819号
学位記番号	甲第550号
学位授与年月日	2017.03.28
学位授与機関	東邦大学
DOI	info:doi/10.1080/14397595.2016.1179860
その他資源識別子	<a href="http://www.tandfonline.com/doi/abs/10.1080/14397595.2016.1179860?journalCode=imor20">http://www.tandfonline.com/doi/abs/10.1080/14397595.2016.1179860?journalCode=imor20</a>
メタデータのURL	<a href="https://mylibrary.toho-u.ac.jp/webopac/TD61681168">https://mylibrary.toho-u.ac.jp/webopac/TD61681168</a>

**Original article****The growth factor midkine may play a pathophysiological role in rheumatoid arthritis**

Emiko Shindo<sup>1</sup>, Toshihiro Nanki<sup>1</sup>, Natsuko Kusunoki<sup>1</sup>, Kotaro Shikano<sup>1</sup>, Mai Kawazoe<sup>1</sup>,  
Hiroshi Sato<sup>1</sup>, Kaichi Kaneko<sup>1</sup>, Sei Muraoka<sup>1</sup>, Makoto Kaburaki<sup>1</sup>, Yoshikiyo Akasaka<sup>2</sup>,  
Hideaki Shimada<sup>3</sup>, Tomoko Hasunuma<sup>1</sup>, and Shinichi Kawai<sup>1</sup>

<sup>1</sup>Division of Rheumatology, Department of Internal Medicine, Toho University School of  
Medicine, 6-11-1, Omori-nishi, Ota-ku, Tokyo, 143-8541 Japan.

<sup>2</sup>Division of Chronic Inflammatory Diseases, Advanced Medical Research Center, Toho  
University Graduate School of Medicine, 6-11-1, Omori-nishi, Ota-ku, Tokyo, 143-8541  
Japan.

<sup>3</sup>Division of General and Gastroenterological Surgery, Department of Surgery, Toho  
University School of Medicine, 6-11-1, Omori-nishi, Ota-ku, Tokyo, 143-8541 Japan.

Number of text pages and figure legends: 20

Number of figures: 6

Number of tables: 0

Keywords: midkine, rheumatoid arthritis, disease activity, chemokine, cytokine

Running title: Midkine is a disease activity marker of RA

Correspondence to:

Toshihiro Nanki, MD, PhD

1 Division of Rheumatology, Department of Internal Medicine,

2 Toho University School of Medicine.

3  
4 6-11-1, Omori-nishi, Ota-ku, Tokyo, 143-8541 Japan.

5  
6  
7 Tel: +81-3-3762-4151, FAX: +81-3-3298-0045

8  
9 E-mail: toshihiro.nanki@med.toho-u.ac.jp  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Abstract**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
*Objectives.* Midkine (MK) is involved in cell proliferation, differentiation, migration, and survival. In this study, we measured serum MK levels in rheumatoid arthritis (RA) and investigated the correlation of serum MK with RA disease activity. Expression and effect of MK in RA synovial tissue were also examined.

12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
*Methods.* Serum MK and production of inflammatory mediators by rheumatoid synovial fibroblasts (RSFs) were measured by enzyme-linked immunosorbent assay. MK expression in synovial tissue was examined by immunohistochemistry. MK receptor expression was analyzed by RT-PCR and Western blotting.

22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
*Results.* RA patients had a significantly higher serum MK level than healthy controls. In RA patients, the MK level was correlated with DAS28-ESR, disability index of the Health Assessment Questionnaire, and rheumatoid factor level. The serum MK level tended to be decreased by anti-TNF therapy. MK was expressed by synovial lining cells in RA synovial tissues and it enhanced the production of IL-6, IL-8 and CCL2 by RSFs. RSFs expressed LDL receptor-related protein 1, candidate receptor for MK.

36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
*Conclusions.* The serum MK level could be a marker of disease activity in RA and an indicator of a poor prognosis. MK may have a role in the pathogenesis of RA via induction of inflammatory mediators.

## Introduction

1  
2 Midkine (MK) is a heparin-binding growth factor that was originally identified as the retinoic  
3 acid-response gene product [1-3]. It has been reported that MK has a role in cell proliferation,  
4 differentiation, migration, and survival [4-6]. In addition, the pathological role of MK in  
5 malignant tumors has been widely analyzed [7-9]. Several types of cancer cells express MK  
6 [1, 10], and cancer patients with high serum levels of MK are reported to have a poor  
7 prognosis [11]. Furthermore, MK enhances tumor cell growth and invasion [12]. Moreover,  
8 MK might modulate inflammatory responses [13, 14], since it has been reported to induce the  
9 migration of inflammatory cells [15-17] and suppresses the expansion of regulatory T cell by  
10 blocking the development of tolerogenic dendritic cells [18]. Thus, MK might have a role in  
11 inflammatory diseases as well as cancer.

12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Rheumatoid arthritis (RA) is characterized by chronic inflammation of multiple joints. The synovial tissue of RA patients features proliferation of synoviocytes, accumulation of inflammatory cells (including lymphocytes and macrophages), production of inflammatory mediators, and angiogenesis [19]. Several biologic agents are effective for RA, including monoclonal antibodies against tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) and interleukin (IL)-6 receptor. However, the risk of severe infection is increased by these biologic agents [20, 21] and some patients do not respond to them. Therefore, novel therapies that are more effective for RA and safer than conventional treatment are still needed.

It has been reported that MK is expressed in RA synovial tissue [15, 22] and that MK levels are elevated in the serum [22] and synovial fluid [15] of RA patients. These reports suggest that MK may play a role in the pathogenesis of RA. In this study, we analyzed the relation between the serum MK level and disease activity in RA, and we also investigated the stimulatory effect of MK on cultured synovial fibroblasts obtained from RA patients.

## Methods

### Samples

Peripheral blood samples were obtained from 146 RA patients and 85 healthy subjects. All of the RA patients fulfilled the classification criteria of the American College of Rheumatology/European League Against Rheumatism [23]. Among the 146 RA patients, 121 (82.8%) were women. The mean age  $\pm$  standard deviation (SD) was  $60.9 \pm 14.0$  years, the disease duration was  $125.7 \pm 115.2$  months, and the C-reactive protein (CRP) level was  $0.57 \pm 0.13$  mg/dl. In addition, the disease activity score (DAS)28-erythrocyte sedimentation rate (ESR) [24] was  $2.76 \pm 1.46$  and the disability index of the Health Assessment Questionnaire (HAQ) [25] was  $0.58 \pm 0.87$ . Of the 146 RA patients, 118 (80.8%) were positive for rheumatoid factor (RF) and 113 (77.4%) were positive for anti-cyclic citrullinated protein (CCP) antibodies. Sixty-eight patients were taking prednisolone and 101 patients were using methotrexate. Other medications were salazosulfapyridine in 2 patients, tacrolimus in 6, cyclosporin in 1, infliximab in 11, etanercept in 28, adalimumab in 7, abatacept in 3, golimumab in 3, and tocilizumab in 13. Blood samples from microscopic polyangiitis (n=3), adult-onset Still's disease (n=2), systemic lupus erythematosus (n=3), and polymyositis (n=2) were also obtained. Among the healthy subjects, 76 (89.4%) were women and their age was  $63.2 \pm 12.1$  years. The ages and gender ratio were not significantly different between RA patients and healthy controls. Blood samples from healthy controls were obtained at Kitasato University Kitasato Institute Hospital. The subjects were healthy on the basis of their medical histories. They did not have a present or past history of organopathy, such as heart, lung, liver or kidney disease. They were normal based on physical examination and clinical laboratory tests. Synovial tissue samples were obtained from 3 patients with RA and 3 patients with osteoarthritis (OA) who underwent total knee or hip replacement. The experimental protocol was approved in advance by the Ethics Committees of Toho Medical Center Omori Hospital

(approval number: 24-207, 26-22), Toho University School of Medicine (2703024007), and Kitasato University Kitasato Institute Hospital (13028).

### **Measurement of MK**

Serum was obtained from peripheral blood samples of the RA, polyangiitis, adult-onset Still's disease, systemic lupus erythematosus, polymyositis and healthy controls and was frozen at -80°C until analysis. Concentrations of MK and RF were determined with enzyme-linked immunosorbent assay (ELISA) kits (Cellmid, Sydney, Australia), according to the manufacturer's instruction. To examine whether RF interfered with the MK assay, serum samples from 3 RA patients were incubated with Immunoglobulin Inhibiting Reagent<sup>®</sup> (Bioreclamation IVT, Baltimore, MD, USA) for 18 h at 4°C before determination of the MK level.

### **Sharp score**

To evaluate radiographic changes of the hands, the Sharp score was calculated [26]. Erosions were evaluated at 16 sites of each hand and wrist and joint space narrowing was assessed at 15 sites on each side. The score for erosions at each site ranged from 0 to 5, giving a total score between 0 and 160, while the score for joint space narrowing at each site ranged from 0 to 4, give a total score between 0 and 120. The score was independently evaluated by ES and KS, and the mean score was calculated.

### **Immunohistochemistry**

Synovial tissues were fixed with freshly prepared 4% (v/v) paraformaldehyde in Tris buffered saline (TBS). Sections (3 µm) were immersed in ethanol containing 3% (v/v) H<sub>2</sub>O<sub>2</sub> for 5 min to block endogenous peroxidase activity. Then the sections were incubated with EzBlock

1 Chemi<sup>®</sup> (Atto, Tokyo, Japan) for 10 min to block nonspecific binding. Sections were  
2 subsequently incubated for 2 h with polyclonal goat anti-human MK antibody (2 µg/ml; R&D  
3 Systems, Minneapolis, MN, USA) or normal goat serum as a control. Next, the sections were  
4 washed four times with TBS and expression was detected using a Vectastain<sup>®</sup> Elite ABC kit  
5 (Vector Laboratories, Burlingame, CA, USA). Finally, the sections were washed in TBS and  
6 counterstained with hematoxylin.  
7  
8  
9  
10  
11  
12

### 13 **Cell culture**

14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
Rheumatoid synovial fibroblasts (RSFs) were prepared from synovial tissue samples as  
described previously [27]. In brief, synovial tissue was digested for 2 h with 0.2% (w/v)  
bacterial collagenase (Wako Pure Chemical Industries, Osaka, Japan) and then suspended in  
Roswell Park Memorial Institute (RPMI) 1640 medium with 10% (v/v) fetal bovine serum  
(FBS), 100 units/ml penicillin, and 100 g/ml streptomycin. The cells were incubated for  
several days at 37°C under 5% CO<sub>2</sub>, after which nonadherent cells were removed.

Fibroblast-like adherent cells from the third or fourth passages were used as RSFs. These  
cells were cultured overnight in 96-well plates (2 x 10<sup>4</sup> cells/well), and then were incubated  
with MK (0, 5, or 50 ng/ml; Wako Pure Chemical Industries) for 48 h at 37°C in RPMI 1640  
medium containing 10% FBS. Levels of IL-6, IL-8 and chemokine (C-C motif) ligand 2  
(CCL2) and chemokine (C-X3-C motif) ligand 1 (CX3CL1) in the culture supernatant were  
assessed by using ELISA kits (Life Technologies, Waltham, MA, USA, for IL-6 and CCL2;  
R&D Systems for IL-8 and CX3CL1), according to the manufacturer's instructions. To  
measure prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) level, RSFs were plated in 96-well plates (2 x 10<sup>4</sup>  
cells/well) and cultured for 48 h at 37°C with MK (0, 5, or 50 ng/ml) in medium containing  
1% FBS. Three µM arachidonic acid (Cayman Chemical, Ann Arbor, MI, USA) was added to  
each well for last 30 min. Then, concentration of PGE<sub>2</sub> in the medium was measured by an



ELISA kit according to the recommendations of the manufacturer (Cayman Chemical).

### Reverse transcription-polymerase chain reaction (RT-PCR)

RSFs were seeded in culture medium containing 10% FBS, and total RNA was extracted with an RNeasy Mini kit according to the recommendations of the manufacturer (QIAGEN GmbH, Hilden, Germany). Reverse transcription was performed with a SuperScript first-strand synthesis system for RT-PCR according to the recommendations of the manufacturer (Invitrogen, Carlsbad, CA, USA), with 1 µg of total RNA from the cells as a template. Equal amounts of each reverse-transcribed product were amplified by PCR with HotStar *Taq* polymerase (QIAGEN GmbH). The primer sequences and numbers of basepairs were as follows: for anaplastic lymphoma kinase (ALK) (814 bp), sense 5'-AGA TCC TCC TGA TGC CCA CT and antisense 5'-AGG ATC TTG TCC TCT CCG CT; for protein tyrosine phosphatase, receptor type Z1 (PTPRZ1) (196 bp), sense 5'-TGC TTC CAG ACT GCA CAC TT and antisense 5'-CCT GGT AAA ACT CTT TCA GTG TCT C; for Notch2 (817 bp), sense 5'-TGG CAC ATG TGT TGA TGG GA and antisense 5'-GTT GGA GAG GCA CTC GTT GA; for LDL receptor-related protein 1 (LRP1) (419 bp), sense 5'-CTG GCG AAC AAA CAC ACT GG and antisense 5'-GCA CGT CAG ACC AGT ACA CA; and for glyceraldehyde-3-phosphate dehydrogenase (GAPDH) (598 bp), sense 5'-CCA CCC ATG GCA AAT TCC ATG GCA and antisense 5'-TCT AGA CGG CAG GTC AGG TCC ACC. After initial denaturation at 95°C for 15 min, PCR involved amplification for 28 cycles of 30 sec at 94°C, 30 sec at 56°C, and 30 sec at 72°C, followed by elongation for 5 min at 72°C. The amplified complementary DNA fragments were resolved by electrophoresis on a 2% agarose gel, and were detected under ultraviolet light using LAS-3000 (Fujifilm, Tokyo, Japan) after staining the gel with ethidium bromide.

## Western blot analysis

1  
2 RSFs were lysed in M-PER<sup>®</sup> mammalian protein extraction reagent (Thermo Fisher  
3  
4 Scientific, Waltham, MA, USA) with Halt<sup>™</sup> protease inhibitor cocktail (Thermo Fisher  
5  
6 Scientific) according to the recommendations of the manufacturer. The protein content of the  
7  
8 lysates was determined with Qubit<sup>®</sup> 3.0 Fluorometer using Qubit<sup>®</sup> protein assay (Thermo  
9  
10 Fisher Scientific). Then cell lysates were adjusted to 30 µg of protein and were applied to  
11  
12 Bolt<sup>®</sup> Bis-Tris Plus gel (4–12%) (Thermo Fisher Scientific) for electrophoresis in MOPS  
13  
14 buffer. Next, the proteins were electroblotted onto Immobilon-P polyvinylidene difluoride  
15  
16 membranes with iBlot<sup>®</sup> 2 Dry Blotting System (Thermo Fisher Scientific). The membranes  
17  
18 were incubated with primary antibodies, monoclonal rabbit anti-human Notch2 antibody  
19  
20 (Cell Signaling Technology, Danvers, MA, USA), monoclonal rabbit anti-human LRP1  
21  
22 antibody (Abcam plc, Cambridge, UK), or monoclonal mouse anti-human GAPDH antibody  
23  
24 (Santa Cruz Biotechnology, Dallas, TX, USA), with a dilution of 1:200 (Notch2), 1:3000  
25  
26 (LRP1) and 1:100 (GAPDH), and the secondary antibody (HRP-conjugated goat anti-rabbit  
27  
28 antibody or HRP-conjugated goat anti-mouse antibody) was added (at a dilution of 1:2000)  
29  
30 and incubation was performed for 2.5 h using iBind Flex Western System (Thermo Fisher  
31  
32 Scientific). Protein bands were detected with enhanced Novex<sup>®</sup> ECL Chemiluminescent  
33  
34 Substrate Reagent Kit (Invitrogen) using LAS-3000 (Fujifilm).  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45

## Statistical analysis

46  
47 Results are expressed as the mean ± SD and/or median with interquartile range. Statistical  
48  
49 analysis was performed with Prism software (ver. 6; GraphPad Software, La Jolla, CA, USA).  
50  
51 The significance of between-group differences of the serum MK concentration was  
52  
53 determined by the Mann-Whitney U-test. Simple linear regression analysis was used to assess  
54  
55 correlations between the serum MK level and the DAS28-ESR, HAQ score, Sharp score, or  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

RF level. To compare IL-6, IL-8, CCL2 and PGE<sub>2</sub> production by RSFs, analysis of variance was applied with Bonferroni's correction. In all analyses,  $p < 0.05$  was considered to indicate statistical significance.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Results

### Serum MK level

As shown in Figure 1, the serum level of MK was significantly higher in the RA patients than in the healthy controls [RA:  $787.7 \pm 68.8$  (mean  $\pm$  SD), 540.8 (391.7-781.1) median (interquartile range); healthy controls:  $373.6 \pm 15.1$ , 334.4 (284.8-426.1) pg/ml,  $p < 0.0001$ ].

We also measured serum MK levels of patients with microscopic polyangiitis, adult-onset Still's disease, systemic lupus erythematosus, and polymyositis. Although the MK concentrations were vary in the patients; microscopic polyangiitis (n=3; 240.5, 458.1, 378.3 pg/ml), adult-onset Still's disease (n=2; 76.7, 232.5), systemic lupus erythematosus (n=3; 146.1, 339.8, 686.6), and polymyositis (n=2; 47.0, 31.0), they were not significantly increased compared to healthy controls. To determine whether the serum MK level could be a useful biomarker of RA disease activity, we examined the correlation between serum MK and DAS28-ESR. In the RA patients, we found that the serum MK level was significantly correlated with DAS28-ESR ( $r=0.223$ ,  $p=0.0194$ ) (Figure 2A). Since HAQ is widely used for measuring functional disability in daily life, the correlation between the HAQ score and serum MK was examined, and a significant correlation of serum MK with the HAQ score was identified ( $r=0.329$ ,  $p=0.0143$ ) (Figure 2B). The Sharp score for radiographic bone destruction of the finger and wrist joints was also analyzed for a correlation with serum MK. There was little correlation between serum MK level and Sharp score; the p value was less than 0.05, but correlation coefficient was less than 0.1 ( $r=0.097$ ,  $p=0.0013$ ) (Figure 2C). Furthermore, serum MK was strongly correlated with the RF titer ( $r=0.615$ ,  $p < 0.00001$ ) (Figure 2D).

Since RF often interferes with the assay other serum proteins, we evaluated the influence of RF on the ELISA for MK. First RF in serum was blocked with Immunoglobulin Inhibiting Reagent<sup>®</sup> and then we measured MK using the ELISA. After blocking RF, the

1 measured concentration of MK was not altered, while the concentration of RF was  
2 significantly decreased (data not shown), suggesting that RF did not affect the assay for MK.  
3  
4 The anti-CCP antibody titer was not correlated with the serum MK level (data not shown).  
5  
6

7 Four of the RA patients started treatment with infliximab, an anti-TNF- $\alpha$  monoclonal  
8 antibody, and the concentration of MK was measured several times. The clinical courses,  
9 including serum MK levels, are shown in Figure 3. In all the patients, CRP and ESR were  
10 lowered after infliximab treatment, and dosage of prednisolone could be decreased. Serum  
11 MK level tended to be lowered although it was sometimes fluctuated.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21

### 22 **MK expression in RA synovium**

23  
24 Expression of MK in the synovial tissues harvested from RA and OA patients was examined  
25 by immunohistochemistry. There was strong expression of MK in the synovial lining cells of  
26 RA synovial tissue (Figure 4), while MK expression was minimal in OA synovial tissue.  
27  
28  
29  
30  
31

### 32 **Stimulatory effect of MK on RSFs**

33  
34 Finally, we investigated the *in vitro* effects of MK on RSFs. After RSFs were incubated with  
35 MK for 48 h, the concentrations of IL-6, IL-8, CCL2, CX3CL1 and PGE<sub>2</sub>, which are thought  
36 to be involved in the pathogenesis of RA [28-30], in the culture supernatant were determined  
37 with ELISA kits. It was found that stimulation with MK significantly increased the  
38 production of IL-6 and IL-8 by RSFs in a dose-dependent manner (Figure 5A and B).  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

66 We also examined the expression of candidate MK receptors, ALK, PTPRZ1,  
67 Notch2 and LRP1 [31], in RSFs. Notch2 and LRP1 mRNAs were expressed, but expression

of ALK and PTPRZ1 was not detected by RT-PCR (Figure 6A). The protein expression of LRP1, but not Notch2, was detected in RSFs by Western blotting (Figure 6B).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Discussion

1  
2 In this study, we showed that the serum MK level is elevated in RA patients and is correlated  
3  
4 with clinical indexes of RA. In addition, MK expression was obvious in RA synovial tissue,  
5  
6 and stimulation of RSFs with MK enhanced the production of IL-6, IL-8 and CCL2.  
7  
8

9  
10 Previous study has shown that the serum MK level is elevated in RA patients and is  
11  
12 correlated with the RF titer [22]. In this study, we found that the serum MK level was  
13  
14 correlated with DAS28-ESR and that MK tended to be decreased after treatment with an  
15  
16 anti-TNF- $\alpha$  antibody. These results indicate that the serum MK level could be a useful marker  
17  
18 of RA disease activity. Moreover, serum MK was correlated with HAQ, indicating that MK  
19  
20 also could be an indicator for a poor prognosis of RA. In fact, MK was also correlated with  
21  
22 the RF titer, which is a well-known poor prognostic factor [32]. Therefore, it is possible that  
23  
24 the serum MK level could be a useful disease activity marker as well as a prognostic factor.  
25  
26  
27 In this study, we retrospectively evaluated clinical courses, including serum MK, ESR and  
28  
29 CRP levels, in 4 patients treated with anti-TNF- $\alpha$  antibody. We further need to prospectively  
30  
31 analyze the changes of serum MK level and the disease activities in detail by the treatment in  
32  
33 large number of patients to identify the characteristics of MK as a marker for activity of RA.  
34  
35  
36  
37

38  
39 We found that MK was expressed in the RA synovial lining, which is consistent with  
40  
41 a previous report [15, 22], while there was minimal MK expression in OA synovium. The  
42  
43 MK level in synovial fluid was also reported to be higher in RA compared with OA [22]. In  
44  
45 the present study, we found that stimulation with MK enhanced the production of IL-6, IL-8  
46  
47 and CCL2 by cultured RSFs. IL-8 is a chemokine (C-X-C motif) ligand 8 (CXCL8), which  
48  
49 induces blood vessel formation and angiogenesis and also exhibits a chemoattractant activity  
50  
51 for neutrophils and dendritic cells [28, 33, 34]. CCL2 is thought to induce migration of  
52  
53 monocytes into inflamed RA synovial tissue [29]. Thus, our findings suggest that MK might  
54  
55 play a role in inflammatory cell migration via enhancement of chemokine production by  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

RSFs. Moreover, it was reported that MK induces neutrophil migration [15, 17], so MK could both directly and indirectly promote inflammatory cell accumulation in RA synovial tissue and synovial fluid. IL-6 has a wide range of effects on immune cells and contributes to the pathogenesis of autoimmune diseases [35], and it has been reported that inhibition of IL-6 signaling by anti-IL-6 or anti-IL-6 receptor monoclonal antibodies is effective for RA [36, 37]. Our findings indicated that MK may contribute to the production of cytokines such as IL-6, which regulate immune cell activation.

As candidate receptors for MK, ALK, PTPRZ1, Notch2, and LRP1 are reported [31]. We found that LPR1 was expressed in mRNA and protein levels in RSFs. LPR1 might be a responsible receptor for MK stimulation in RSFs.

It has been reported that MK enhances endothelial cell proliferation and vascular growth [12] and also upregulates the differentiation of osteoclasts *in vitro* [22]. Therefore, MK may contribute to angiogenesis and bone destruction in the synovial tissues in RA, as well as stimulating RSFs. In a murine model of antibody-induced arthritis, it was found that MK expression was upregulated in the affected joints after arthritis developed [22]. In addition, development of antibody-induced arthritis was significantly inhibited in MK-deficient mice compared with wild-type mice [22]. Thus, MK could be a potential therapeutic target in patients with RA.

In conclusion, the serum MK level could be a marker of RA disease activity and an indicator of a poor prognosis. It also seems possible that MK plays a role in the pathogenesis of RA via induction of IL-6, IL-8 and CCL2.



## Acknowledgements

This work was supported in part by a Project Research Grant (26-25, 27-12) from Toho University School of Medicine to ES; Heisei 23 & 26 Strategic Research Foundation Grants from the Japanese Ministry of Education, Culture, Sports, Science & Technology to YA and SK; and a Grant from Japan Society for the Promotion of Science (23591449) to SK. We wish to thank Ms. Sonoko Sakurai for secretarial assistance.

## Competing interests

Hideaki Shimada received research grant from Cellmid Ltd. Other authors declare that they have no competing interests.

**References**

1. Muramatsu T. Midkine and pleiotrophin: two related proteins involved in development, survival, inflammation and tumorigenesis. *J Biochem.* 2002; 132: 359-71.
2. Muramatsu T. Midkine (MK), the product of a retinoic acid responsive gene, and pleiotrophin constitute a new protein family regulating growth and differentiation. *Int J Dev Biol.* 1993; 37: 183-8.
3. Kadomatsu K, Muramatsu T. Midkine and pleiotrophin in neural development and cancer. *Cancer Lett.* 2004; 204: 127-43.
4. Muramatsu H, Shirahama H, Yonezawa S, Maruta H, Muramatsu T. Midkine, a retinoic acid-inducible growth/differentiation factor: immunochemical evidence for the function and distribution. *Dev Biol.* 1993; 159: 392-402.
5. Owada K, Sanjo N, Kobayashi T, Mizusawa H, Muramatsu H, Muramatsu T, et al. Midkine inhibits caspase-dependent apoptosis via the activation of mitogen-activated protein kinase and phosphatidylinositol 3-kinase in cultured neurons. *J Neurochem.* 1999; 73: 2084-92.
6. Maeda N, Ichihara-Tanaka K, Kimura T, Kadomatsu K, Muramatsu T, Noda M. A receptor-like protein-tyrosine phosphatase PTPzeta/RPTPbeta binds a heparin-binding growth factor midkine. Involvement of arginine 78 of midkine in the high affinity binding to PTPzeta. *J Biol Chem.* 1999; 274: 12474-9.
7. Aridome K, Tsutsui J, Takao S, Kadomatsu K, Ozawa M, Aikou T, et al. Increased midkine gene expression in human gastrointestinal cancers. *Jpn J Cancer Res.* 1995; 86: 655-61.
8. Koide N, Hada H, Shinji T, Ujike K, Hirasaki S, Yumoto Y, et al. Expression of the midkine gene in human hepatocellular carcinomas. *Hepatogastroenterology.* 1999; 46: 3189-96.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
9. Konishi N, Nakamura M, Nakaoka S, Hiasa Y, Cho M, Uemura H, et al. Immunohistochemical analysis of midkine expression in human prostate carcinoma. *Oncology*. 1999; 57: 253-7.
10. Tsutsui J, Kadomatsu K, Matsubara S, Nakagawara A, Hamanoue M, Takao S, et al. A new family of heparin-binding growth/differentiation factors: increased midkine expression in Wilms' tumor and other human carcinomas. *Cancer Res*. 1993; 53: 1281-5.
11. Ikematsu S, Nakagawara A, Nakamura Y, Ohira M, Shinjo M, Kishida S, et al. Plasma midkine level is a prognostic factor for human neuroblastoma. *Cancer Sci*. 2008; 99: 2070-4.
12. Choudhuri R, Zhang HT, Donnini S, Ziche M, Bicknell R. An angiogenic role for the neurokines midkine and pleiotrophin in tumorigenesis. *Cancer Res*. 1997; 57: 1814-9.
13. Muramatsu T. Midkine, a heparin-binding cytokine with multiple roles in development, repair and diseases. *Proc Jpn Acad Ser B Phys Biol Sci*. 2010; 86: 410-25.
14. Weckbach LT, Muramatsu T, Walzog B. Midkine in inflammation. *ScientificWorldJournal*. 2011; 11: 2491-505.
15. Takada T, Toriyama K, Muramatsu H, Song XJ, Torii S, Muramatsu T. Midkine, a retinoic acid-inducible heparin-binding cytokine in inflammatory responses: chemotactic activity to neutrophils and association with inflammatory synovitis. *J Biochem*. 1997; 122: 453-8.
16. Horiba M, Kadomatsu K, Nakamura E, Muramatsu H, Ikematsu S, Sakuma S, et al. Neointima formation in a restenosis model is suppressed in midkine-deficient mice. *J Clin Invest*. 2000; 105: 489-95.
17. Sato W, Kadomatsu K, Yuzawa Y, Muramatsu H, Hotta N, Matsuo S, et al. Midkine is involved in neutrophil infiltration into the tubulointerstitium in ischemic renal injury. *J Immunol*. 2001; 167: 3463-9.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
18. Sonobe Y, Li H, Jin S, Kishida S, Kadomatsu K, Takeuchi H, et al. Midkine inhibits inducible regulatory T cell differentiation by suppressing the development of tolerogenic dendritic cells. *J Immunol.* 2012; 188: 2602-11.
19. Feldmann M, Brennan FM, Maini RN. Role of cytokines in rheumatoid arthritis. *Annu Rev Immunol.* 1996; 14: 397-440.
20. Komano Y, Tanaka M, Nanki T, Koike R, Sakai R, Kameda H, et al. Incidence and risk factors for serious infection in patients with rheumatoid arthritis treated with tumor necrosis factor inhibitors: a report from the Registry of Japanese Rheumatoid Arthritis Patients for Longterm Safety. *J Rheumatol.* 2011; 38: 1258-64.
21. Sakai R, Cho SK, Nanki T, Watanabe K, Yamazaki H, Tanaka M, et al. Head-to-head comparison of the safety of tocilizumab and tumor necrosis factor inhibitors in rheumatoid arthritis patients (RA) in clinical practice: results from the registry of Japanese RA patients on biologics for long-term safety (REAL) registry. *Arthritis Res Ther.* 2015; 17: 74.
22. Maruyama K, Muramatsu H, Ishiguro N, Muramatsu T. Midkine, a heparin-binding growth factor, is fundamentally involved in the pathogenesis of rheumatoid arthritis. *Arthritis Rheum.* 2004; 50: 1420-9.
23. Aletaha D, Neogi T, Silman AJ, Funovits J, Felson DT, Bingham CO, 3rd, et al. 2010 rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Ann Rheum Dis.* 2010; 69: 1580-8.
24. Felson DT, Smolen JS, Wells G, Zhang B, van Tuyl LH, Funovits J, et al. American College of Rheumatology/European League against Rheumatism provisional definition of remission in rheumatoid arthritis for clinical trials. *Ann Rheum Dis.* 2011; 70: 404-13.
25. Fries JF. The hierarchy of quality-of-life assessment, the Health Assessment

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
- Questionnaire (HAQ), and issues mandating development of a toxicity index. *Clin Trials*. 1991; 12: 106S-117S.
26. van der Heijde DM, van Riel PL, Nuver-Zwart IH, Gribnau FW, van de Putte LB. Effects of hydroxychloroquine and sulphasalazine on progression of joint damage in rheumatoid arthritis. *Lancet*. 1989; 333: 1036-8.
27. Kusunoki N, Yamazaki R, Kawai S. Induction of apoptosis in rheumatoid synovial fibroblasts by celecoxib, but not by other selective cyclooxygenase 2 inhibitors. *Arthritis Rheum*. 2002; 46: 3159-67.
28. Ritchlin C. Fibroblast biology. Effector signals released by the synovial fibroblast in arthritis. *Arthritis Res*. 2000; 2: 356-60.
29. Hayashida K, Nanki T, Girschick H, Yavuz S, Ochi T, Lipsky PE. Synovial stromal cells from rheumatoid arthritis patients attract monocytes by producing MCP-1 and IL-8. *Arthritis Res*. 2001; 3: 118-26.
30. Nanki T, Imai T, Nagasaka K, Urasaki Y, Nonomura Y, Taniguchi K, et al. Migration of CX3CR1-positive T cells producing type 1 cytokines and cytotoxic molecules into the synovium of patients with rheumatoid arthritis. *Arthritis Rheum*. 2002; 46: 2878-83.
31. Kadomatsu K, Kishida S, Tsubota S. The heparin-binding growth factor midkine: the biological activities and candidate receptors. *J Biochem*. 2013; 153: 511-21.
32. Vencovsky J, Machacek S, Sedova L, Kafkova J, Gatterova J, Pesakova V, et al. Autoantibodies can be prognostic markers of an erosive disease in early rheumatoid arthritis. *Ann Rheum Dis*. 2003; 62: 427-30.
33. Murphy PM, Tiffany HL. Cloning of complementary DNA encoding a functional human interleukin-8 receptor. *Science*. 1991. 253: 1280-3.
34. Rubbert A, Combadiere C, Ostrowski M, Arthos J, Dybul M, Machado E, et al. Dendritic cells express multiple chemokine receptors used as coreceptors for HIV entry. *J Immunol*.

1998; 160: 3933-41.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
35. Tanaka T, Narazaki M, Kishimoto T. Therapeutic targeting of the interleukin-6 receptor. *Annu Rev Pharmacol Toxicol.* 2012; 52: 199-219.
36. Maini RN, Taylor PC, Szechinski J, Pavelka K, Broll J, Balint G, et al. Double-blind randomized controlled clinical trial of the interleukin-6 receptor antagonist, tocilizumab, in European patients with rheumatoid arthritis who had an incomplete response to methotrexate. *Arthritis Rheum.* 2006; 54: 2817-929.
37. Mease P, Strand V, Shalamberidze L, Dimic A, Raskina T, Xu LA, et al. A phase II, double-blind, randomised, placebo-controlled study of BMS945429 (ALD518) in patients with rheumatoid arthritis with an inadequate response to methotrexate. *Ann Rheum Dis.* 2012; 71: 1183-9.

## Figure legends

1  
2 Figure 1. Serum MK level. The serum MK concentration was determined by ELISA in 146  
3  
4 RA patients and 85 healthy controls.  
5  
6

7  
8  
9 Figure 2. Correlation of the serum MK level with the DAS28-ESR, HAQ, Sharp score, and  
10  
11 RF in the patients with RA. Correlations between MK and the DAS28-ESR (A), HAQ score  
12  
13 (B), Sharp score (C), and RF (D) were determined in the 146 patients. Each point represents  
14  
15 an individual patient.  $P < 0.05$  was considered to indicate statistical significance.  
16  
17  
18  
19  
20

21  
22 Figure 3. Clinical courses by anti-TNF- $\alpha$  treatment. Clinical courses of each individual,  
23  
24 including the treatment, serum levels of MK, ESR and CRP in 4 patients who received  
25  
26 infliximab, are shown (A to D). PSL: prednisolone, MTX: methotrexate, IFX: infliximab.  
27  
28  
29  
30

31  
32 Figure 4. Expression of MK in synovial tissue. MK expression in synovial tissue harvested  
33  
34 from RA patients (n=3) and OA patients (n=3) was analyzed by immunohistochemistry.  
35  
36 Sections were counterstained with hematoxylin and representative photomicrographs are  
37  
38 shown. Bar =100  $\mu\text{m}$ . MK was strongly expressed by synovial lining cells in RA synovial  
39  
40 tissue, whereas MK expression was minimal in OA synovial tissue.  
41  
42  
43  
44  
45

46  
47 Figure 5. Stimulatory effect of MK on RSFs. After RSFs were incubated with MK for 48 h,  
48  
49 the concentrations of IL-6 (A), IL-8 (B), CCL2 (C) and PGE<sub>2</sub> (D) in the culture supernatant  
50  
51 were measured by ELISA. Data are the mean  $\pm$  SD for one of three independent experiments  
52  
53 analyzed in triplicate. \* $p < 0.05$ , \*\* $p < 0.005$  vs. MK (0 ng/ml).  
54  
55  
56  
57

58  
59 Figure 6. Expression of candidate receptors for MK in RSFs. Expression of mRNA of ALK,  
60  
61  
62  
63  
64  
65

PTPRZ1, Notch2, and LRP1 was analyzed by RT-PCR in RSFs from 3 RA patients (A).

Expression of Notch2 and LRP1 in 3 RSFs was examined by Western blotting (B). M:

molecular weight marker.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65






Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6