

東邦大学学術リポジトリ

Toho University Academic Repository

タイトル	Molecular characterization of Neisseria gonorrhoeae isolates collected through a national surveillance programme in Japan, 2013: Evidence of the emergence of a ceftriaxone resistant strain from a ceftriaxonesusceptible lineage
別タイトル	2013年に日本の全国サーベイランスプログラムで分離された淋菌の分子疫学解析:セフトリアキソン感性系統からセフトリアキソン耐性株が出現した科学的根拠
作成者(著者)	綾見(花尾), 麻美
公開者	東邦大学
発行日	2022.03.16
掲載情報	東邦大学大学院医学研究科 博士論文. 8.
資料種別	学位論文
内容記述	主査: 中島耕一 / タイトル: Molecular characterization of Neisseria gonorrhoeae isolates collected through a national surveillance programme in Japan, 2013: Evidence of the emergence of a ceftriaxone resistant strain from a ceftriaxonesusceptible lineage / 著者: Mami Hanao, Kotaro Aoki, Yoshikazu Ishii, Ken Shimuta, Makoto Ohnishi, Kazuhiro Tateda / 掲載誌: Journal of Antimicrobial Chemotherapy / 巻号・発行年等: 76(7): 1769-1775, 2021 / 本文ファイル: 査読後原稿
著者版フラグ	ETD
報告番号	32661甲第1021号
学位記番号	甲第700号
学位授与年月日	2022.03.16
学位授与機関	東邦大学
DOI	info:doi/10.1093/jac/dkab104
その他資源識別子	https://academic.oup.com/jac/article/76/7/1769/6261155
メタデータのURL	https://mylibrary.toho-u.ac.jp/webopac/TD78930228

Molecular characterization of *Neisseria gonorrhoeae* isolates collected through a national surveillance programme in Japan, 2013: evidence of the emergence of a ceftriaxone-resistant strain from a ceftriaxone-susceptible lineage

Mami HANA^{1,2}, Kotaro AOKI^{3*}, Yoshikazu ISHII^{1,3}, Ken SHIMUTA^{4,5}, Makoto OHNISHI⁴ and Kazuhiro TATEDA^{1,3}

¹Department of Microbiology and Infectious Diseases, Toho University Graduate School of Medicine, Tokyo, Japan; ²Department of Medical Technology, Faculty of Health Science, Tokyo University of Technology, Tokyo, Japan; ³Department of Microbiology and Infectious Diseases, Toho University School of Medicine, Tokyo, Japan; ⁴Department of Bacteriology I, National Institute of Infectious Diseases, Tokyo, Japan; ⁵Antimicrobial Resistance Research Center, National Institute of Infectious Diseases, Tokyo, Japan

Running title: Molecular characterization of *N. gonorrhoeae* in Japan, 2013

*Corresponding author. E-mail: kotaro.aoki@med.toho-u.ac.jp

Abstract

Objectives: To investigate the spread of ceftriaxone-resistant *Neisseria gonorrhoeae* lineages similar to strains H041 (2009) and FC428 (2015), we characterized 55 strains collected in 2013 from hospitals across Japan.

Methods: Susceptibility testing and whole-genome sequencing.

Results: Susceptibility rates were 58% for cefixime and 98% for ceftriaxone. The 55 strains were whole-genome sequenced and classified into nine MLST-STs. MLST-ST1901 was the most prevalent ($n=19$) followed by MLST-ST7363 ($n=12$) and MLST-ST7359 ($n=11$). The most prevalent *penA* [encoding penicillin binding protein 2 (PBP2)] mosaic types, based on the *N. gonorrhoeae* sequence typing for antimicrobial resistance (NG-STAR) scheme, were 10.001 ($n=20$) followed by 34.001 ($n=13$). The H041 and FC428 strains were not detected; however, a single ceftriaxone-resistant strain (TUM15748) with a MIC of 0.5 mg/L ceftriaxone was identified. The TUM15748 strain belonged to MLST-ST7359 and *N. gonorrhoeae* multiantigen sequence typing-ST6771, and had a novel PBP2 (PBP2_{TUM15748}, *penA* type 169.001). The amino acid sequence of PBP2_{TUM15748} showed partial similarity to that of PBP2 from *N. gonorrhoeae* GU140106 and commensal *Neisseria perflava* and *Neisseria cinerea*. Natural transformation and recombination experiments using full-length TUM15748 *penA* showed that the ceftriaxone MICs of transformants increased 16-fold or more compared with the parental ceftriaxone-susceptible recipient strain (NG9807, belonging to MLST-ST7363). No ceftriaxone-resistant MLST-ST7359 strains have previously been reported.

Conclusions: We showed here that a ceftriaxone-susceptible lineage acquired a mutant PBP2 mosaic type, integrating partial PBP2 sequences from commensal *Neisseria* species, resulting in the emergence of ceftriaxone-resistant strains.

Introduction

Resistance of *Neisseria gonorrhoeae* to extended-spectrum cephalosporins (ESCs) is a global public health concern.^{1,2} Ceftriaxone is one of the last remaining recommended therapies for treatment of gonococcal infections. Thus, dissemination of ceftriaxone-resistant isolates may limit the availability of medical interventions.³⁻⁵

The main determinants of resistance to ESCs are mosaic *penA* alleles encoding penicillin-binding protein 2 (PBP2) variants with reduced affinity for ESCs.⁶ *N. gonorrhoeae* can import extracellular DNA fragment by natural transformation and homologous recombination, thus acquiring antimicrobial resistance determinants from other *Neisseria* species.⁷ Transformation of mosaic *penA* alleles can increase the MIC of ceftriaxone for recipient cells.⁸⁻¹⁰ In Japan, the first high-level ceftriaxone-resistant gonococcal strain (H041) was isolated in Kyoto in 2009.⁸ Subsequently, intensified surveillance conducted from 2010 to 2012 in the same district did not identify dissemination of the H041 strain.¹¹ Five years after the emergence of H041, two ceftriaxone-resistant strains belonging to different lineages (GU140106¹² and FC428¹³) were isolated in Japan (in 2014 and 2015, respectively). Their mosaic *penA* alleles were proposed to originate from commensal *Neisseria* species and parts of the *penA* sequences of GU140106 and FC428 were similar to *Neisseria cinerea penA*.¹³⁻¹⁶

Worldwide, the most prevalent MLST-STs of cefixime- and ceftriaxone-resistant gonococcal strains are MLST-ST7363, MLST-ST1901, and MLST-ST1903.^{13,17-20} The ceftriaxone-resistant

H041,⁸ A8806,²¹ and GU140106¹² strains belong to MLST-ST7363, the F89 strain²² belongs to MLST-ST1901, and the FC428 strain¹³ belongs to MLST-ST1903. Most *N. gonorrhoeae* strains with mosaic *penA* alleles were classified as MLST-ST1901 and MLST-ST7363, and therefore close attention must be paid to trends in the ceftriaxone resistance of strains belonging to these two lineages.^{13,17} However, the diversity of mosaic *penA* alleles is not always reflected in MLST results because *penA* and housekeeping genes can transform separately. A full explanation of the characteristics of resistant strains requires whole-genome sequences.

Genomic data on ESC-susceptible and -resistant *N. gonorrhoeae* clinical strains in Japan are limited.²³ Only a few reports have described the sequences of isolates from outpatient clinics in the area where H041 was detected (Kyoto and Osaka, Japan), from 2010 to 2012 and in 2015.^{11,13}

In this study, we characterized 55 *N. gonorrhoeae* strains collected through national surveillance efforts conducted 4 years following the detection of the high-level ceftriaxone-resistant H041 and 2 years prior to detection of the ceftriaxone-resistant FC428 strain. We determined the draft whole-genome sequences of these strains and analysed the mechanisms underlying the development of ceftriaxone resistance by *in vitro* natural transformation of *penA* alleles.

Materials and methods

Strains

In 2013, 55 *N. gonorrhoeae* clinical isolates were identified at 25 medical facilities in 16 prefectures in Japan as part of a post-marketing survey of levofloxacin.²⁴ This study was conducted with approval of the Ethics Committee of the Faculty of Medicine, Toho University (no. A16039).

Antimicrobial susceptibility testing

MICs were determined by the agar dilution method according to the CLSI M07-ED11 guidelines.²⁵ The results were interpreted according to the European Committee on Antimicrobial Susceptibility Testing (EUCAST) clinical breakpoints version 10.0 (https://www.eucast.org/clinical_breakpoints/). Because the breakpoint for levofloxacin is indeterminate, that of ciprofloxacin was substituted. Strains were incubated on GC agar for 24 h at 35°C in the presence of 5% CO₂. The MICs of the following six antimicrobial agents were measured: penicillin G (concentration range: 0.063–16 mg/L), cefixime (0.063–16 mg/L), ceftriaxone (0.063–4 mg/L), spectinomycin (0.063–64 mg/L), levofloxacin (0.063–64 mg/L) and azithromycin (0.063–64 mg/L). *N. gonorrhoeae* strains ATCC 49226, NG9807⁸ and FC428¹³ were used for antimicrobial susceptibility testing quality control.

Molecular characterization by draft whole-genome sequencing

Genomic DNA was obtained by phenol/chloroform extraction of bacteria and purified using a QIAamp PCR purification kit (Qiagen, Valencia, CA, USA). To determine draft whole-genome

sequences, DNA libraries were prepared using the Nextera XT DNA library preparation kit (Illumina, San Diego, CA, USA). Libraries were sequenced on a MiSeq system using a 600-cycle reagent kit (2×300 bp paired-end reads) and draft genome contigs were obtained by *de novo* assembly using SPAdes (v3.13.0). MLST and *N. gonorrhoeae* multiantigen sequence typing (NG-MAST) were performed using the pubMLST (<https://pubmlst.org/neisseria>)²⁶ and NG-MAST (<https://www.ng-mast.net>)²⁷ schemes, respectively. Genotyping of *penA* (encoding PBP2) was conducted using the *N. gonorrhoeae* sequence typing for antimicrobial resistance (NG-STAR) scheme (<https://ngstar.canada.ca>).²⁸ Multiple sequence alignment used CLASTALW and phylogenetic analyses used Maximum Likelihood were performed using MEGA X.²⁹ Core genome SNP-based phylogenetic analysis was performed using draft whole genome sequences. The MiSeq sequencing data were aligned to the genome sequence of the reference *N. gonorrhoeae* strain WHO Y (GenBank accession no. LT592161) using the Burrows-Wheeler Aligner (BWA) with ‘MEM’ option.³⁰ We constructed a core genome alignment using SAMtools (version 1.1) mpileup³¹ and VarScan (version 2.3.7) mpileup2cns³² and a maximum-likelihood phylogenetic tree using PhyML.³³ Using this as the starting tree, we inferred homologous recombination events that imported DNA fragments from outside the ST and constructed a clonal phylogeny with corrected branch lengths using ClonalFrameML.³⁴ A final phylogenetic tree was constructed using RAxML (version 8.2.12).³⁵

***penA* natural transformation**

To assess whether the *penA* gene (encoding PBP2) of the ceftriaxone-resistant strain TUM15748 (*penA*_{TUM15748}) contributed to the increased MIC of ceftriaxone, full-length *penA*_{TUM15748} was amplified by PCR and a recipient strain, *N. gonorrhoeae* NG9807, was transformed with the *penA*_{TUM15748} amplicon as previously described.³⁶ *N. gonorrhoeae* NG9807, which belongs to MLST-ST7363, NG-MAST-ST4093, had a ceftriaxone MIC of 0.016 mg/L.

Complete whole-genome sequencing

To identify any unintended mutations and recombination events in the transformant strain chromosomes, we determined the complete whole-genome sequences of the donor, recipient, and transformant strains. High molecular weight DNA was extracted using NucleoBond AXG 20 (TaKaRa, Shiga, Japan) and prepared for MinION (Oxford Nanopore Technologies, Oxford, UK) sequencing. DNA libraries were prepared using the Rapid Barcoding Kit (Oxford Nanopore Technologies). Pooled libraries were sequenced using a Nanopore flow cell R9.4. Hybrid *de novo* assemblies of Illumina and MinION reads were performed with Unicycler (v0.4.8-beta).³⁷ Additional nucleotide error correction was performed using pilon (v1.22) three times.³⁸ Gene annotation and analysis of genome structure was performed using DFAST (v1.1.15) (<https://dfast.nig.ac.jp/>) and MAUVE (v2.1.1),³⁹ respectively.

Nucleotide sequence accession numbers

The sequences determined in this study were deposited in the DNA Data Bank of Japan (DDBJ) under BioProject number PRJDB9563. The draft genome sequences of 55 *N. gonorrhoeae* strains were deposited in the DDBJ and under GenBank accession numbers SAMD00217291 to SAMD00217345. The accession numbers of the complete genome sequences determined in this study were SAMD00220608 for TUM15748, SAMD00220610 for TUM16691 (NG9807), and SAMD00257605 for TUM19855 (NG9807/PBP2_{TUM15748}). Datasets S1 and S2, available as Supplementary data at *JAC* Online, show the sequencing data statistics.

Results

Antimicrobial susceptibility

The results of the antimicrobial susceptibility testing of the 55 *N. gonorrhoeae* isolates are summarized in Table 1 and Figure 1. Of the 55 isolates, 42% were susceptible to cefixime and 98% were susceptible to ceftriaxone. The susceptibility rates to benzylpenicillin, azithromycin and levofloxacin were 2%, 95% and 25%, respectively. All 55 strains were susceptible to spectinomycin. Only one strain, *N. gonorrhoeae* TUM15748, was resistant to ceftriaxone (MIC: 0.5 mg/L). This strain had low susceptibility to benzylpenicillin (MIC: 0.5 mg/L) but was susceptible to azithromycin, spectinomycin and levofloxacin (Table 1).

Genotyping and phylogenetic analysis using MLST, NG-MAST and core genome SNPs

A total of 55 draft whole-genome sequences were obtained at an average depth of 51.0-fold (SD 18.8-fold) (see Dataset S1). Assembled genomes had an average of 218.0 (SD 119.0) contigs and an average N50 value of 24683.0 (SD 7095.0) bp. Genotyping of the 55 *N. gonorrhoeae* strains by MLST and NG-MAST is summarized in Table S1. The 55 strains were classified into nine MLST-STs. MLST-ST1901 was the most prevalent ($n=19$, 35%), followed by MLST-ST7363 ($n=12$, 22%) and MLST-ST7359 ($n=11$, 20%). The 55 strains were classified into 38 NG-MAST-STs. NG-MAST-ST6543, NG-MAST-ST6771 and NG-MAST-ST6798 were the most prevalent ($n=4$, 7% each), followed by NG-MAST-ST1407 ($n=3$, 5%). Each NG-MAST-ST could be assigned to a specific MLST-ST, except for NG-MAST-ST4018, which was associated with both MLST-ST1594 and MLST-ST15378. MLST-ST15378, NG-MAST-ST19619, NG-MAST-ST19620, NG-MAST-ST19621, NG-MAST-ST19622, NG-MAST-ST19623, and NG-MAST-ST19706 have not been reported previously. Of the 55 *N. gonorrhoeae* strains, three shared the same STs as the ceftriaxone-resistant strain F89 (also called strain WHO Y) (MLST-ST1901 and NG-MAST-ST1407)²² and four shared the same STs with the ceftriaxone-resistant strain GU140106 (MLST-ST7363 and NG-MAST-ST6543).¹²

Core genome SNP-based phylogeny, antimicrobial susceptibility, and determinants of antimicrobial resistance

The relationships between a core genome SNP-based phylogeny, antimicrobial susceptibility, and antimicrobial resistance determinants are shown in Figure 1. Of the 55 strains, 53 could be divided into three clades (A, B, and C). Clades A, B, and C could be subdivided into two (A-1 and A-2), three (B-1, B-2, and B-3), and two (C-1 and C-2) subclades, respectively. The two remaining isolates (TUM15789 and TUM15795) were not closely related to other strains and were independent of each other. Three MLST-STs (MLST-ST7359, MLST-ST1594, and MLST-ST15378) belonged to clade A. Except for TUM15748, subclade A-2 consisted of ESC-susceptible *N. gonorrhoeae* strains. The cefixime- and ceftriaxone-resistant strain TUM15748 harbored a novel *penA* type 169 (mosaic) (Figure 1, Figure 2, and Table S1). Subclade B-1 consisted of MLST-ST1901 and MLST-ST7360. Subclades B-2 and B-3 were mixed, containing MLST-ST1901 and MLST-ST1579. Subclade B-2 harboured strains of *penA* types 34.001 (mosaic) and 72.001 (mosaic). Except for TUM15759, subclade B-3 harboured strains of *penA* type 10.001 (mosaic). All clade B strains harboured mutations in GyrA (S91F and D95G) and ParC (S87R) and were resistant to levofloxacin. Clade C consisted of MLST-ST7363. Subclade C-1 harboured strains of *penA* types 10.001 (mosaic) and 168.001 (mosaic). Subclade C-2 harboured strains of *penA* type 150.001 (semi-mosaic). Subclade C-1 strains showed reduced susceptibility to cefixime. Of the three subclade C-2 *penA* type 150.001 strains, only one showed reduced susceptibility to cefixime. In clade C, only one *penA* type 10.001 strain and one *penA* type 168.001 strain showed reduced susceptibility to ceftriaxone. All clade C strains harbouring mutations in GyrA (S91F and D95N) and ParC (S87R and S88P) were resistant to levofloxacin. All

subclade A-2 strains harboured the MtrR A39T mutation and nearly all clade B strains harboured the -35A deletion. Almost all clade B and all subclade C-1 strains harbored the PorB G120K and A121D mutations. No penicillinase genes or 23S rRNA gene mutations were detected in any strains.

Characteristics of the PBP2_{TUM15748} amino acid sequence

Phylogenetic analysis was performed to investigate the relevance of the novel PBP2 sequence of TUM15748 (PBP2_{TUM15748}) through comparison with previously reported PBP2 sequences (Figure 2). The amino acid sequence of PBP2_{TUM15748} was most similar to that of PBP2_{GU140106} (94.7% identity). PBP2_{TUM15748} bore two conserved amino acid substitutions, A311V and T483S, which contribute to reduced susceptibility to ESCs.^{8,40} Regional similarity of PBP2 amino acid sequences across *Neisseria* species is shown in Figure 3. Based on regional amino acid sequence similarity, PBP2 could be divided into seven regions. PBP2_{TUM15748} was highly similar (>95% identity) to the region-1, -2, -5, and -7 sequences of PBP2_{GU140106}. PBP2_{TUM15748} shared high similarity in region-1, -2, -3, and -5 with *N. perflava* PBP2_{NSU-per_52} and region-2, -3, -5, and -7 of *N. cinerea* PBP_{SI57-1}. Interestingly, the sequences of these four PBP2s were highly similar in region-5, and except for PBP2_{NSU-per_52}, shared a conserved A311V amino acid substitution.

PBP2_{TUM15748} raised the ESC MICs of susceptible strains

Ceftriaxone-susceptible *N. gonorrhoeae* NG9807 was transformed with *penA*_{TUM15748}. The ceftriaxone and cefixime MICs of transformant NG9807/PBP2_{TUM15748} increased 16-fold or more compared with the recipient strain (Table 2). By contrast, the MIC of benzylpenicillin was unaffected. The complete whole-genome sequence of the donor, recipient, and transformant strains showed that the recipient full-length *penA* was completely homologous with *penA*_{TUM15748}. No unexpected mutations or recombination events were detected in transformant chromosomes.

Discussion

Following the identification of the high-level ceftriaxone-resistant *N. gonorrhoeae* H041 strain in 2009, the spread of this strain has been carefully monitored through surveillance efforts in Japan.^{11,41–43} Fortunately, the H041 strain has not been detected, and we also did not observe dissemination of H041 in this study. The ceftriaxone-resistant FC428 strain has likewise not been detected. The rate of azithromycin susceptibility in the present study (95%) was higher than that reported in a previous study (76.7%) undertaken in Japan from 2010 to 2012 ($n=193$).¹¹ Almost all *N. gonorrhoeae* strains with reduced susceptibility to ESCs identified to date have belonged to MLST-ST1901, MLST-ST1903, or MLST-ST7363.^{8,13,22,43,44} In this study, we found that of 32 gonococcal strains with reduced susceptibility or resistance to cefixime, 31 belonged to clade B and subclade C-1 according to core genome SNP-based phylogenetic analysis. Almost all strains belonging to clade B and subclade C-1 were MLST-ST1901, MLST-ST7363, and MLST-ST1579. The mutations G120K and A121D in PorB have been reported to produce slight reductions in

susceptibility to penicillin, cefixime, and ceftriaxone,^{45,46} and therefore these mutations contribute to the rising MIC baselines of penicillin, cefixime, and ceftriaxone in most strains of clade B and subclade C-1 (Figure 1). Our data showed that the MLST-ST1579 lineage may be developing reduced susceptibility to cefixime in Japan; only one strain of ceftriaxone-resistant *N. gonorrhoeae* of MLST-ST1579 has been previously reported.⁴⁷ Assessment of quinolone- and macrolide-resistance determinants, as well as PBP2 sequences, using draft whole-genome sequences is important to understand the molecular epidemiology of antimicrobial resistance in *N. gonorrhoeae*.

In this study, we found that the ESC-susceptible lineage MLST-ST7359 included only one ceftriaxone-resistant strain, TUM15748. According to previous studies in Japan, all MLST-ST7359 strains were susceptible to both cefixime and ceftriaxone.^{11,43} TUM15748 was inferred to have developed ceftriaxone resistance by acquiring a novel type 169.001 *penA* (mosaic) encoding a PBP2 variant with decreased affinity for ceftriaxone. PBP2_{TUM15748} shared regional amino acid similarity with PBP2_{SI57-1} and PBP2_{NSU-per_52} from commensal *N. cinerea* and *N. perflava*, respectively (Figure 3). Our data support the possibility that the previously reported partial sequence of *N. cinerea* PBP2 is a component of PBP2 variants with decreased ceftriaxone affinity in *N. gonorrhoeae*.¹⁶ There has been no report of strains with genetic similarity to TUM15748 since TUM15748 was isolated. Mutations in *penA* are predicted to have a negative impact on gonococcal fitness,^{8,48} so it is possible that PBP2_{TUM15748} imposes fitness costs on *N. gonorrhoeae*, limiting its spread.

The number of samples analysed in our study was quite small compared with the estimated number of gonorrhoea cases in 2013 in Japan (25 606). Therefore, low sample size is one possible reason for our inability to detect strain H041 in this study. However, we detected the ceftriaxone-resistant *N.*

gonorrhoeae strain TUM15748 harbouring a novel *penA* allele. This finding suggests that novel *penA* alleles associated with reduced susceptibility to ESCs may have a prevalence of around 2%.

Appropriate use of ESCs is required to avoid selecting for ESC-resistant *N. gonorrhoeae* strains emerging at low frequency.

In this study, we did not detect the high-level ceftriaxone-resistant H041 strain and the ceftriaxone-resistant FC428 strain in 2013 in Japan. One *N. gonorrhoeae* strain in the ESC-susceptible lineage MLST-ST7359 acquired resistance to ceftriaxone mediated by a novel *penA* allele.

Acknowledgements

We thank the Levofloxacin Surveillance group for providing *N. gonorrhoeae* strains and Daiichi Sankyo Company, Ltd, for their support of the study. We thank Edanz Group (<https://en-author-services.edanzgroup.com/ac>) for editing a draft of this manuscript.

Funding

This study was supported by the Japan Agency for Medical Research and Development (AMED) under Grant Number JP19fk0108062h0302.

Transparency declarations

None to declare.

Author contributions

M.H. and K.A. were co-investigators and responsible for the data analysis. Y.I. was responsible for the organization and coordination of the study. K.S. and M.O. provided research strains and guided some experiments. K.T. was the primary investigator. All authors contributed to the writing of the final manuscript and met the ICMJE authorship criteria.

Supplementary data

Table S1 and Datasets S1 and S2 are available as Supplementary data at *JAC* Online.

References

1 Unemo M, Lahra MM, Cole M *et al.* World Health Organization Global Gonococcal Antimicrobial Surveillance Program (WHO GASP): Review of new data and evidence to inform international collaborative actions and research efforts. *Sex Health* 2019; **16**: 412–25.

2 CDC. *Antibiotic resistance threats in the United States*. 2019; 1–150.

<https://www.cdc.gov/drugresistance/pdf/threats-report/2019-ar-threats-report-508.pdf>.

3 Fifer H, Natarajan U, Alexander S *et al*. Failure of dual antimicrobial therapy in treatment of gonorrhoea. *N Engl J Med* 2016; **374**: 2502–4.

4 Town K, Obi C, Quaye N *et al*. Drifting towards ceftriaxone treatment failure in gonorrhoea: Risk factor analysis of data from the Gonococcal Resistance to Antimicrobials Surveillance Programme in England and Wales. *Sex Transm Infect* 2017; **93**: 39–45.

5 Chen MY, Stevens K, Tideman R *et al*. Failure of 500 mg of ceftriaxone to eradicate pharyngeal gonorrhoea, Australia. *J Antimicrob Chemother* 2013; **68**: 1445–7.

6 Unemo M, Shafer WM. Antimicrobial resistance in *Neisseria gonorrhoeae* in the 21st Century: Past, evolution, and future. *Clin Microbiol Rev* 2014; **27**: 587–613.

7 Hamilton HL, Dillard JP. Natural transformation of *Neisseria gonorrhoeae*: From DNA donation to homologous recombination. *Mol Microbiol* 2006; **59**: 376–85.

8 Ohnishi M, Golparian D, Shimuta K *et al*. Is *Neisseria gonorrhoeae* initiating a future era of untreatable gonorrhoea?: Detailed characterization of the first strain with high-level resistance to ceftriaxone. *Antimicrob Agents Chemother* 2011; **55**: 3538–45.

- 9** Lindberg R, Fredlund H, Nicholas R *et al.* *Neisseria gonorrhoeae* isolates with reduced susceptibility to cefixime and ceftriaxone: Association with genetic polymorphisms in *penA*, *mtrR*, *porB1b*, and *ponA*. *Antimicrob Agents Chemother* 2007; **51**: 2117–22.
- 10** Zhao S, Duncan M, Tomberg J *et al.* Genetics of chromosomally mediated intermediate resistance to ceftriaxone and cefixime in *Neisseria gonorrhoeae*. *Antimicrob Agents Chemother* 2009; **53**: 3744–51.
- 11** Shimuta K, Unemo M, Nakayama SI *et al.* Antimicrobial resistance and molecular typing of *Neisseria gonorrhoeae* isolates in Kyoto and Osaka, Japan, 2010 to 2012: Intensified surveillance after identification of the first strain (H041) with high-level ceftriaxone resistance. *Antimicrob Agents Chemother* 2013; **57**: 5225–32.
- 12** Deguchi T, Yasuda M, Hatazaki K *et al.* New clinical strain of *Neisseria gonorrhoeae* with decreased susceptibility to ceftriaxone, Japan. *Emerg Infect Dis* 2016; **22**: 142–4.
- 13** Nakayama S, Shimuta K, Furubayashi K *et al.* New ceftriaxone- and multidrug-resistant *Neisseria gonorrhoeae* strain with a novel mosaic *penA* gene isolated in Japan. *Antimicrob Agents Chemother* 2016; **60**: 4339–41.
- 14** Ameyama S, Onodera S, Takahata M *et al.* Mosaic-like structure of penicillin-binding protein 2 gene (*penA*) in clinical isolates of *Neisseria gonorrhoeae* with reduced susceptibility to cefixime. *Antimicrob Agents Chemother* 2002; **46**: 3744–9.

- 15** Ito M, Deguchi T, Mizutani KS *et al.* Emergence and spread of *Neisseria gonorrhoeae* clinical isolates harboring mosaic-like structure of penicillin-binding protein 2 in central Japan. *Antimicrob Agents Chemother* 2005; **49**: 137-143.
- 16** Igawa G, Yamagishi Y, Lee KI *et al.* *Neisseria cinerea* with high ceftriaxone MIC is a source of ceftriaxone and cefixime resistance-mediating *penA* sequences in *Neisseria gonorrhoeae*. *Antimicrob Agents Chemother* 2018; **62**: 1–5.
- 17** Lahra MM, Martin I, Demczuk W *et al.* Cooperative recognition of internationally disseminated ceftriaxone-resistant *Neisseria gonorrhoeae* strain. *Emerg Infect Dis* 2018; **24**: 735–40.
- 18** Unemo M, Nicholas RA. Emergence of multidrug-resistant, extensively drug-resistant and untreatable gonorrhoea. *Future Microbiol* 2012; **7**: 1401–22.
- 19** Unemo M, Golparian D, Sánchez-Busó L *et al.* The novel 2016 WHO *Neisseria gonorrhoeae* reference strains for global quality assurance of laboratory investigations: Phenotypic, genetic and reference genome characterization. *J Antimicrob Chemother* 2016; **71**: 3096–108.
- 20** Lee K, Nakayama S, Osawa K *et al.* Clonal expansion and spread of the ceftriaxone-resistant *Neisseria gonorrhoeae* strain FC428, identified in Japan in 2015, and closely related isolates. *J Antimicrob Chemother* 2019; **74**: 1812–9.
- 21** Lahra MM, Ryder N, Whiley DM. A new multidrug-resistant strain of *Neisseria gonorrhoeae* in Australia. *N Engl J Med* 2014; **371**: 1849–50.

22 Unemo M, Golparian D, Nicholas R *et al.* High-level cefixime- and ceftriaxone-resistant *Neisseria gonorrhoeae* in France: Novel *penA* mosaic allele in a successful international clone causes treatment failure. *Antimicrob Agents Chemother* 2012; **56**: 1273–80.

23 Yahara K, Nakayama S, Shimuta K *et al.* Genomic surveillance of *Neisseria gonorrhoeae* to investigate the distribution and evolution of antimicrobial-resistance determinants and lineages. *Microb Genomics* 2018; **4**: 1–3.

24 Yamaguchi K, Ohno A, Ishii Y *et al.* Surveillance of in vitro susceptibilities to levofloxacin and various antibacterial agents for 11,762 clinical isolates obtained from 69 centers in 2013. *Jpn J Antibiot* 2016; **69**: 1–25.

25 CLSI. *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically, 11th Edition* 2018.

26 Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics: BIGSdb software, the PubMLST.org website and their applications [version 1; referees: 2 approved]. *Wellcome Open Res* 2018; **3**: 1–20.

27 Martin IMC, Ison CA, Aanensen DM *et al.* Rapid sequence-based identification of gonococcal transmission clusters in a large metropolitan area. *J Infect Dis* 2004; **189**: 1497–505.

- 28** Demczuk W, Sidhu S, Unemo M *et al.* *Neisseria gonorrhoeae* sequence typing for antimicrobial resistance, a novel antimicrobial resistance multilocus typing scheme for tracking global dissemination of *N. gonorrhoeae* strains. *J Clin Microbiol* 2017; **55**: 1454–68.
- 29** Kumar S, Stecher G, Li M *et al.* MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol* 2018; **35**: 1547–9.
- 30** Li H, Durbin R. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* 2010; **26**: 589–95.
- 31** Li H, Handsaker B, Wysoker A *et al.* The sequence alignment/map format and SAMtools. *Bioinformatics* 2009; **25**: 2078–9.
- 32** Koboldt DC, Zhang Q, Larson DE *et al.* VarScan 2: Somatic mutation and copy number alteration discovery in cancer by exome sequencing. *Genome Res* 2012; **22**: 568–76.
- 33** Guindon S, Dufayard JF, Lefort V *et al.* New algorithms and methods to estimate maximum-likelihood phylogenies: Assessing the performance of PhyML 3.0. *Syst Biol* 2010; **59**: 307–21.
- 34** Didelot X, Wilson DJ. ClonalFrameML: Efficient inference of recombination in whole bacterial genomes. *PLoS Comput Biol* 2015; **11**: 1–18.
- 35** Stamatakis A. RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 2014; **30**: 1312–3.

- 36** Ohnishi M, Watanabe Y, Ono E *et al.* Spread of a chromosomal cefixime-resistant *penA* gene among different *Neisseria gonorrhoeae* lineages. *Antimicrob Agents Chemother* 2010; **54**: 1060–7.
- 37** Wick RR, Judd LM, Gorrie CL *et al.* Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads. *PLOS Comput Biol* 2017; **13**: e1005595.
- 38** Walker BJ, Abeel T, Shea T *et al.* Pilon: An integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One* 2014; **9**: e112963.
- 39** Darling ACE, Mau B, Blattner FR *et al.* Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res* 2004; **14**: 1394–403.
- 40** Tomberg J, Unemo M, Ohnishi M *et al.* Identification of amino acids conferring high-level resistance to expanded-spectrum cephalosporins in the *penA* gene from *Neisseria gonorrhoeae* Strain H041. *Antimicrob Agents Chemother* 2013; **57**: 3029–36.
- 41** Hamasuna R, Yasuda M, Ishikawa K *et al.* Nationwide surveillance of the antimicrobial susceptibility of *Neisseria gonorrhoeae* from male urethritis in Japan. *J Infect Chemother* 2013; **19**: 571–8.
- 42** Hamasuna R, Yasuda M, Ishikawa K *et al.* The second nationwide surveillance of the antimicrobial susceptibility of *Neisseria gonorrhoeae* from male urethritis in Japan, 2012–2013. *J Infect Chemother* 2015; **21**: 340–5.

- 43** Shimuta K, Watanabe Y, Nakayama S *et al.* Emergence and evolution of internationally disseminated cephalosporin-resistant *Neisseria gonorrhoeae* clones from 1995 to 2005 in Japan. *BMC Infect Dis* 2015; **15**: 378.
- 44** Chen SC, Yuan LF, Zhu XY *et al.* Sustained transmission of the ceftriaxone-resistant *Neisseria gonorrhoeae* FC428 clone in China. *J Antimicrob Chemother* 2020; **75**: 2499–502.
- 45** Olesky M, Hobbs M, Nicholas RA. Identification and analysis of amino acid mutations in porin IB that mediate intermediate-level resistance to penicillin and tetracycline in *Neisseria gonorrhoeae*. *Antimicrob Agents Chemother* 2002; **46**: 2811–20.
- 46** Parmar NR, Singh R, Martin I *et al.* Genomic analysis reveals antibiotic-susceptible clones and emerging resistance in *Neisseria gonorrhoeae* in Saskatchewan, Canada. *Antimicrob Agents Chemother* 2020; **64**: 1–7.
- 47** de Curraize C, Kumanski S, Micaëlo M *et al.* Ceftriaxone-resistant *Neisseria gonorrhoeae* Isolates (2010 to 2014) in France characterized by using whole-genome sequencing. *Antimicrob Agents Chemother* 2016; **60**: 6962–4.
- 48** Vincent LR, Kerr SR, Tan Y *et al.* In vivo -selected compensatory mutations restore the fitness cost of mosaic *penA* alleles that confer ceftriaxone resistance in *Neisseria gonorrhoeae*. *MBio* 2018; **9**: 1–18.

49 Shimuta K, Igawa G, Yasuda M *et al.* A real-time PCR assay for detecting a *penA* mutation associated with ceftriaxone resistance in *Neisseria gonorrhoeae*. *J Glob Antimicrob Resist* 2019; **19**: 46–9.

Table 1. Antimicrobial susceptibility of 55 *Neisseria gonorrhoeae* strains identified in a national hospital surveillance study in Japan, 2013

Antimicrobial	Breakpoints ^a (mg/L)	No. (%) of isolates ^d			MIC ₅₀ ^e	MIC ₉₀ ^f
		S	I	R		
Benzylpenicillin	S ≤0.06/R >1	1 (2)	31 (56)	23 (42)	1	4
Cefixime	S ≤0.125/R >0.125	23 (42)	0 (0)	32 (58)	0.25	0.5
Ceftriaxone	S ≤0.125/R >0.125	54 (98)	0 (0)	1 (2)	0.06	0.125
Azithromycin	ECOFF 1 mg/L ^b	52 (95)	0 (0)	3 (5)	0.25	0.5
Spectinomycin	S ≤64/R >64	55 (100)	0 (0)	0 (0)	16	16
Levofloxacin	S ≤0.03/R >0.06 ^c	14 (25)	0 (0)	41 (75)	8	16

^aEuropean Committee on Antimicrobial Susceptibility Testing breakpoints version 10.0 are shown.

^bThe clinical breakpoint for azithromycin monotherapy is undefined. With the aim of detecting acquired resistance mechanisms, the ECOFF (epidemiological cut-off value) is 1 mg/L. Susceptible strains have MICs below the ECOFF and resistant strains have MICs equal to or greater than the ECOFF.

^cThe breakpoint of ciprofloxacin was applied to levofloxacin.

^dS, susceptible; I, intermediate; R, resistant.

^eMIC₅₀, MIC of an antimicrobial inhibiting 50% of isolates.

^fMIC₉₀, MIC of an antimicrobial inhibiting 90% of isolates.

Table 2. Phylogenetic and genetic characteristics of *N. gonorrhoeae* strains used for the transformation assay

Characteristic	Donor	Recipient	Transformant
Strain name	TUM15748 (this study)	NG9807 (9)	NG9807/PBP2 _{TUM15748}
MLST-ST	7359	7363	7363
NG-MAST-ST	6771	4093	4093
<i>penA</i> type	169.001 ^a	2.001	169.001
MIC (mg/L)			
Benzylpenicillin	0.5	2	2
Cefixime	2	<0.06	4
Ceftriaxone	0.5	<0.06	0.5

Abbreviations: NG-MAST-ST, *Neisseria gonorrhoeae* multiantigen sequence typing-defined

sequence type; PBP2, penicillin binding protein 2.

^aNew *penA* type identified in this study.

Figure legends


Figure 1. Phylogenetic relationships, antimicrobial susceptibility, and antimicrobial resistance determinants of *N. gonorrhoeae* strains analysed in this study. Phylogenetic analysis was performed using core-genome single-nucleotide polymorphisms and strain STs were characterized by MLST. Genotyping was conducted using the *N. gonorrhoeae* sequence typing for antimicrobial resistance (NG-STAR) scheme. The MICs of penicillin G (PCG), cefixime (CFIX), ceftriaxone (CTRX), levofloxacin (LVFX), azithromycin (AZM) and spectinomycin (SPCT) were determined. S, susceptible; I, intermediate; R, resistant. The scale distance corresponds to the number of substitutions per site. This figure appears in colour in the online version of *JAC* and in black and white in the print version of *JAC*.

Figure 2. Phylogenetic analysis of PBP2 amino acid sequences determined this study and from previously reported *Neisseria gonorrhoeae* strains and commensal *Neisseria* spp.⁴⁹. This phylogenetic tree was constructed using maximum likelihood methods in MEGA X. Sequence accession numbers were as follows: B904096.1, *Neisseria perflava* NSU-per_52 PBP2; LC410041, *Neisseria polysaccharea* SI28-1 PBP2; LC410042, *Neisseria lactamica* SI60-1 PBP2; LC410043, *Neisseria cinerea* SI57-1 PBP2; LC410044, *Neisseria subflava* SI94-3 PBP2, AB904142, *N. subflava* SH43-1 PBP2; AB904147, *Neisseria* sp. SH43-3 PBP2; AB904125, *N. lactamica* NLA-1 PBP2; AB904126, *N. perflava* NSU-per_57 PBP2; and LC316656, *N. cinerea* AM1601 PBP2. PBP2_{TUM15748} is highlighted in grey. Commensal *Neisseria* spp. are shown in bold text. The scale distance corresponds to the number of substitutions per site.

Figure 3. Regional amino acid sequence similarity of PBP2. PBP2_{TUM15748} amino acid sequence similarity with PBP2_{GU140106}, PBP2_{NSU-per_52}, and PBP2_{SI57-1} was calculated by splitting PBP2 into 50 amino acid segments and sliding the 50-residue window by one amino acid at a time. PBP2-encoding *penA* sequences were downloaded from NG-STAR (<https://ngstar.canada.ca/alleles/penA>). *Neisseria*

perflava NSU-per_52 and *Neisseria cinerea* SI57-1 PBP2 amino acid sequences were downloaded from GenBank (accession nos. BAO96787.1 and LC410043, respectively). This figure appears in colour in the online version of *JAC* and in black and white in the print version of *JAC*.

Figure 1.



Legend

Objects	Value	Color
MLST-ST	1579	
	1594	
	1597	
	1901	
	7359	
	7360	
	7363	
	7827	
	15378	
	1.001 ⁿ	
2.002 ⁿ		
5.002 ⁿ		
10.001 ^m		
34.001 ^m		
72.001 ^m		
106.001 ⁿ		
150.001 sm		
168.001 ^m		
169.001 ^m		
PBP1 (<i>penA</i>) WT	L421P	
	WT	
GyrA	WT	
	S91F, D95G	
	S91F, D95N	
ParC	WT	
	D86N	
	S87R	
23S rRNA	S87R, S88P	
	WT	
MtrR	WT	
	A39T	
	-35A del	
PorB	WT	
	G120K, A121D	
MIC (mg/L)	<=0.06	
	0.125	
	0.25	
	0.5	
	1	
	2	
4		
8		
16		
32		

Figure 2.




Figure 3.

