

東邦大学学術リポジトリ

Toho University Academic Repository

タイトル	Erratum to: Preoperative Chemotherapy, Previous Deep Vein Thrombosis/Pulmonary Thromboembolism, and Old age are Predictors of Preoperative Deep Vein Thrombosis
作成者（著者）	Takayuki, Sugano / Masashi, Uzawa / Kenichiro, Koda / Megumi, Tagami / Takayuki, Kitamura
公開者	The Medical Society of Toho University
発行日	2018.06.01
ISSN	21891990
掲載情報	Toho Journal of Medicine. 4(2). p.81 81.
資料種別	学術雑誌論文
内容記述	ERRATUM
著者版フラグ	publisher
DOI	info:doi/10.14994/tohojmed.2017 3 4 116
メタデータのURL	https://mylibrary.toho u.ac.jp/webopac/TD82514099

Erratum

Erratum to: Preoperative Chemotherapy, Previous Deep Vein Thrombosis/Pulmonary Thromboembolism, and Old age are Predictors of Preoperative Deep Vein Thrombosis

Takayuki Sugano* Masashi Uzawa Kenichiro Koda
Megumi Tagami and Takayuki Kitamura

Department of Anesthesiology (Sakura), Toho University School of Medicine

The original article was published in Toho Journal of Medicine 2017 3 (4): 116-124.

The original version of this article unfortunately contained errors in Table 4. The correct table should be shown as follows:

Table 4 Presence of DVT by each risk factor

	DVT (+) group n = 31	DVT (-) group n = 476	Multivariate analysis		p-value
			Odds ratio	95% CI	
Sex					
Male	15 (48.4%)	203 (42.6%)	0.576	0.224-1.48	0.254
Female	16 (51.6%)	273 (57.4%)			
Age (years)	71 ± 8.38	53 ± 17.12	1.08	1.04-1.13	3.83 × 10 ⁻⁴ *
BMI (kg/m ²)	21.8 ± 3.13	22.8 ± 3.90	0.966	0.862-1.08	0.55
Immobility	4 (12.9%)	20 (4.2%)	0.864	0.151-4.94	0.869
Lower extremity paresis	2 (6.5%)	3 (0.6%)	4.12	0.332-51.1	0.271
Cancer	13 (41.9%)	131 (27.5%)	0.857	0.296-2.48	0.775
Cancer therapy	8 (25.8%)	12 (2.5%)	9.76	2.39-39.9	0.0015 *
Venous compression	0 (0%)	20 (4.2%)	2.05 × 10 ⁻⁷	0-Inf	0.995
Previous DVT/PTE	10 (32.3%)	3 (0.6%)	30.6	6.70-140	1.02 × 10 ⁻⁵ *
Pregnancy and the postpartum period	0 (0%)	18 (3.8%)	7.76 × 10 ⁻⁷	0-Inf	0.996
Estrogen-containing oral contraceptives or hormone replacement therapy	0 (0%)	17 (3.6%)	2.76 × 10 ⁻⁷	0-Inf	0.995
Inflammatory bowel disease	1 (3.2%)	20 (4.2%)	3.56	0.38-33.5	0.266

Numbers of cases are shown with percentages in brackets. Values are expressed as means ± standard deviation for continuous variables.

DVT: deep vein thrombus, PTE: pulmonary thromboembolism, DVT (+) group: patients with actual thrombi, DVT (-) group: patients with D-dimer ≤ 0.72 μg/mL (*n* = 326) and patients in whom no thrombi were detected by lower limb ultrasonography (*n* = 150), BMI: body mass index, Inf: infinity, *: *p* < 0.05

The online version of the original article can be found under doi: 10.14994/tohojmed.2017-3-4-116

564-1 Shimoshizu, Sakura, Chiba 285-8741, Japan

*Corresponding Author: tel: +81-(0)43-462-8811

e-mail: t-sugano@sakura.med.toho-u.ac.jp

DOI: 10.14994/tohojmed.2017-3-4-116

Received Dec. 2, 2016; Accepted Oct. 20, 2017

Toho Journal of Medicine 3 (4), Dec. 1, 2017.

ISSN 2189-1990, CODEN: TJMOA2