

東邦大学学術リポジトリ



OPAC

東邦大学メディアセンター

| | |
|-----------|---|
| タイトル | Erratum to: Preoperative Chemotherapy, Previous Deep Vein Thrombosis/Pulmonary Thromboembolism, and Old age are Predictors of Preoperative Deep Vein Thrombosis |
| 作成者（著者） | Takayuki, Sugano / Masashi, Uzawa / Kenichiro, Koda / Megumi, Tagami / Takayuki, Kitamura |
| 公開者 | The Medical Society of Toho University |
| 発行日 | 2018.06.01 |
| ISSN | 21891990 |
| 掲載情報 | Toho Journal of Medicine. 4(2). p.81 81. |
| 資料種別 | 学術雑誌論文 |
| 内容記述 | ERRATUM |
| 著者版フラグ | publisher |
| DOI | info:doi/10.14994/tohojmed.2017 3 4 116 |
| メタデータのURL | https://mylibrary.toho u.ac.jp/webopac/TD82514099 |

Erratum

Erratum to: Preoperative Chemotherapy, Previous Deep Vein Thrombosis/Pulmonary Thromboembolism, and Old age are Predictors of Preoperative Deep Vein Thrombosis

Takayuki Sugano* Masashi Uzawa Kenichiro Koda
Megumi Tagami and Takayuki Kitamura

Department of Anesthesiology (Sakura), Toho University School of Medicine

The original article was published in Toho Journal of Medicine 2017 3 (4): 116-124.

The original version of this article unfortunately contained errors in Table 4. The correct table should be shown as follows:

Table 4 Presence of DVT by each risk factor

| | DVT (+) group n = 31 | DVT (-) group n = 476 | Multivariate analysis | | p-value |
|---|-------------------------|--------------------------|-------------------------|------------|---------------------------|
| | | | Odds ratio | 95% CI | |
| Sex | | | | | |
| Male | 15 (48.4%) | 203 (42.6%) | 0.576 | 0.224-1.48 | 0.254 |
| Female | 16 (51.6%) | 273 (57.4%) | | | |
| Age (years) | 71 ± 8.38 | 53 ± 17.12 | 1.08 | 1.04-1.13 | 3.83 × 10 ⁻⁴ * |
| BMI (kg/m ²) | 21.8 ± 3.13 | 22.8 ± 3.90 | 0.966 | 0.862-1.08 | 0.55 |
| Immobility | 4 (12.9%) | 20 (4.2%) | 0.864 | 0.151-4.94 | 0.869 |
| Lower extremity paresis | 2 (6.5%) | 3 (0.6%) | 4.12 | 0.332-51.1 | 0.271 |
| Cancer | 13 (41.9%) | 131 (27.5%) | 0.857 | 0.296-2.48 | 0.775 |
| Cancer therapy | 8 (25.8%) | 12 (2.5%) | 9.76 | 2.39-39.9 | 0.0015 * |
| Venous compression | 0 (0%) | 20 (4.2%) | 2.05 × 10 ⁻⁷ | 0-Inf | 0.995 |
| Previous DVT/PTE | 10 (32.3%) | 3 (0.6%) | 30.6 | 6.70-140 | 1.02 × 10 ⁻⁵ * |
| Pregnancy and the postpartum period | 0 (0%) | 18 (3.8%) | 7.76 × 10 ⁻⁷ | 0-Inf | 0.996 |
| Estrogen-containing oral contraceptives or hormone replacement therapy | 0 (0%) | 17 (3.6%) | 2.76 × 10 ⁻⁷ | 0-Inf | 0.995 |
| Inflammatory bowel disease | 1 (3.2%) | 20 (4.2%) | 3.56 | 0.38-33.5 | 0.266 |

Numbers of cases are shown with percentages in brackets. Values are expressed as means ± standard deviation for continuous variables.

DVT: deep vein thrombus, PTE: pulmonary thromboembolism, DVT (+) group: patients with actual thrombi, DVT (-) group: patients with D-dimer ≤ 0.72 μg/mL (*n* = 326) and patients in whom no thrombi were detected by lower limb ultrasonography (*n* = 150), BMI: body mass index, Inf: infinity, *: *p* < 0.05

The online version of the original article can be found under doi: 10.14994/tohojmed.2017-3-4-116

564-1 Shimoshizu, Sakura, Chiba 285-8741, Japan

*Corresponding Author: tel: +81-(0)43-462-8811

e-mail: t-sugano@sakura.med.toho-u.ac.jp

DOI: 10.14994/tohojmed.2017-3-4-116

Received Dec. 2, 2016; Accepted Oct. 20, 2017

Toho Journal of Medicine 3 (4), Dec. 1, 2017.

ISSN 2189-1990, CODEN: TJMOA2